

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|--|---|
| (51) International Patent Classification ⁶: C07D 241/38, 403/04, 401/12, C07C 233/54, 247/14, 233/43, 233/56, A61K 31/495 | A1 | (11) International Publication Number: WO 96/17832 (43) International Publication Date: 13 June 1996 (13.06.96) |
| (21) International Application Number: PCT/US95/14571 (22) International Filing Date: 7 November 1995 (07.11.95) (30) Priority Data: 08/350,765 7 December 1994 (07.12.94) US 08/534,526 23 October 1995 (23.10.95) US (71) Applicant: WARNER-LAMBERT COMPANY [US/US]; 210 Tabor Road, Morris Plains, NJ 07950 (US). (72) Inventors: BIGGE, Christopher, Franklin; 1811 Packard, Ann Arbor, MI 48104 (US). RETZ, Daniel, Martin; 1493 Ainsley Street, Ypsilanti, MI 48197 (US). (74) Agents: RYAN, M., Andrea; Warner-Lambert Company, 201 Tabor Road, Morris Plains, NJ 07950 (US) et al. | (81) Designated States: AU, CA, CZ, EE, HU, JP, LT, LV, MX, NZ, PL, RO, RU, SI, SK, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> | |

(54) Title: NOVEL GLUTAMATE RECEPTOR ANTAGONISTS: FUSED CYCLOALKYLQUINOXALINEDIONES**(57) Abstract**

Novel substituted alicyclic fused ring 2,3-quinoxalinediones, pharmaceutical compositions containing the same and the method of using the same, for the blockade of glutamate receptors, including either or both N-methyl-D-aspartate (NMDA) receptors and non-NMDA receptors such as the α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) receptor and the kainate receptor, are described. Methods of preparing the substituted alicyclic fused ring 2,3-quinoxalinediones are also described. Novel intermediates of the inventive quinoxalinediones are disclosed. The novel substituted alicyclic fused ring 2,3-quinoxalinediones may be used, for example, as neuroprotective agents, for treatment of chronic neurodegenerative disorders, as anticonvulsants and in the treatment of schizophrenia, epilepsy, anxiety, pain and drug addiction.

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|---------------------------------------|----|--------------------------|
| AT | Austria | GB | United Kingdom | MR | Mauritania |
| AU | Australia | GE | Georgia | MW | Malawi |
| BB | Barbados | GN | Guinea | NE | Niger |
| BE | Belgium | GR | Greece | NL | Netherlands |
| BF | Burkina Faso | HU | Hungary | NO | Norway |
| BG | Bulgaria | IE | Ireland | NZ | New Zealand |
| BJ | Benin | IT | Italy | PL | Poland |
| BR | Brazil | JP | Japan | PT | Portugal |
| BY | Belarus | KE | Kenya | RO | Romania |
| CA | Canada | KG | Kyrgyzstan | RU | Russian Federation |
| CF | Central African Republic | KP | Democratic People's Republic of Korea | SD | Sudan |
| CG | Congo | KR | Republic of Korea | SE | Sweden |
| CH | Switzerland | KZ | Kazakhstan | SI | Slovenia |
| CI | Côte d'Ivoire | LJ | Liechtenstein | SK | Slovakia |
| CM | Cameroon | LK | Sri Lanka | SN | Senegal |
| CN | China | LU | Luxembourg | TD | Chad |
| CS | Czechoslovakia | LV | Latvia | TG | Togo |
| CZ | Czech Republic | MC | Monaco | TJ | Tajikistan |
| DE | Germany | MD | Republic of Moldova | TT | Trinidad and Tobago |
| DK | Denmark | MG | Madagascar | UA | Ukraine |
| ES | Spain | ML | Mali | US | United States of America |
| FI | Finland | MN | Mongolia | UZ | Uzbekistan |
| FR | France | | | VN | Viet Nam |
| GA | Gabon | | | | |

- 1 -

TITLE

NOVEL GLUTAMATE RECEPTOR ANTAGONISTS:
FUSED CYCLOALKYLOUINOXALINEDIONES

5

BACKGROUND OF THE INVENTIONField of the Invention

10

This invention is related to 2,3-quinoxalinediones having a substituted alicyclic ring fused with the quinoxaline system. The substituted alicyclic ring fused 2,3-quinoxalinediones are active as excitatory amino acid receptor antagonists acting at glutamate receptors, including either or both N-methyl-D-aspartate (NMDA) receptors and non-NMDA receptors such as the α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) receptor and the kainate receptor. The invention also relates to the use of those quinoxalinediones as neuroprotective agents for treating conditions such as cerebral ischemia or cerebral infarction resulting from a range of

- 2 -


phenomena, such as thromboembolic or hemorrhagic stroke, cerebral vasospasms, hypoglycemia, cardiac arrest, status epilepticus, perinatal asphyxia, anoxia such as from drowning, pulmonary surgery and cerebral trauma, as well as to treat chronic neurodegenerative disorders such as lathyrism, Alzheimer's Disease, Parkinsonism and Huntington's Disease and as anticonvulsants. The compounds of the present invention may also be useful in the treatment of schizophrenia, epilepsy, anxiety, pain and drug addiction.

Related Background Art

Excessive excitation by neurotransmitters can cause the degeneration and death of neurons. It is believed that this degeneration is in part mediated by the excitotoxic actions of the excitatory amino acids (EAA) glutamate and aspartate at the N-methyl-D-aspartate (NMDA) receptor, the α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) receptor, and the kainate receptor. AMPA/kainate receptors may be referred to jointly as non-NMDA receptors. This excitotoxic action is considered responsible for the loss of neurons in cerebrovascular disorders such as cerebral ischemia or cerebral infarction resulting from a range of conditions, such as thromboembolic or hemorrhagic stroke, cerebral vasospasms, hypoglycemia, cardiac arrest, status epilepticus, perinatal asphyxia, anoxia such as from drowning, pulmonary surgery and cerebral trauma, as well as lathyrism, Alzheimer's Disease, Parkinson's Disease and Huntington's Disease.

Several classes of quinoxalinedione derivatives have been disclosed as glutamate (EAA) receptor antagonists. For example, U.S. Patent No. 4,889,855, generically discloses compounds of the formula:

- 3 -



wherein R₁, R₂ and R₃ are independently H, halogen, CN, NH₂, NO₂, SO₃H, SO₂NH₂, and CONH₂. This reference
5 specifically discloses 6-amino, 6-cyano, 5-carbamoyl, 6-nitro and 5,6-dinitro-7,8,9,10-tetrahydro-2,3-dihydroxybenzo(f)quinoxalines. The reference, however, does not disclose any compounds with substitution of the fused cyclohexyl ring i.e., where R₃ is other than
10 hydrogen. Nor does the reference disclose or suggest any methods which would allow substitution of the cyclohexyl ring. U.S. Patent No. 5,081,123 and U.S. Patent No. 5,308,845, describe similar structures except that there is respectively a hydroxy or alkoxy
15 function at the nitrogen on the quinoxalinedione skeleton. Again, however, these references do not suggest or illustrate any examples of substituted cyclohexyl rings.

20 Having both NMDA and non-NMDA antagonist properties in a single entity may provide a superior pharmacological profile. Combinations of NMDA and non-NMDA receptor antagonists have shown synergistic activity in focal and global ischemia [K. Lippert, M. Welsch and J.
25 Krieglstein, Eur. J. Pharmacol. 253 (3), 207-13 (1994)], as anticonvulsants [W. Loescher, C. Rundfelt, D. Hoenack, Eur. J. Neurosci. 5 (11), 1545-50 (1993)], and in protection of neuronal degeneration in retina [J. Mosinger, M. Price, H. Bai, H. Xiao, D. Wozniak and
30 J. Olney, Exp. Neurol. 113, 10-17 (1991)].

Among excitatory amino acid receptor antagonists recognized for usefulness in the treatment of disorders

- 4 -

are those that block AMPA receptors [C.F. Bigge and T.C. Malone, Curr. Opin. Ther. Pat., 951 (1993); M. A. Rogawski, TIPS 14, 325 (1993)]. AMPA receptor antagonists have prevented neuronal injury in several models of global cerebral ischemia [H. Li and A. M. Buchan, J. Cerebr. Blood Flow Metab. 13, 933 (1993); B. Nellgård and T. Wieloch, J. Cerebr. Blood Flow Metab. 12, 2 (1992)] and focal cerebral ischemia [R. Bullock, D. I. Graham, S. Swanson, J. McCulloch, J. Cerebr. Blood Flow Metab. 14, 466 (1994); D. Xue, Z.-G. Huang, K. Barnes, H. J. Lesiuk, K. E. Smith, A. M. Buchan, J. Cerebr. Blood Flow Metab. 14, 251 (1994)]. AMPA antagonists have also shown efficacy in models for analgesia [X.-J. Xu, J.-X. Hao, A. Seiger, Z. Wiesenfeld-Hallin, J. Pharmacol. Exp. Ther. 267, 140 (1993)], and epilepsy [T. Namba, K. Morimoto, K. Sato, N. Yamada, S. Kuroda, Brain Res. 638, 36 (1994); S. E. Brown, J. McCulloch, Brain Res. 641, 10 (1994); S. I. Yamaguchi, S. D. Donevan, M. A. Rogawski, Epilepsy Res. 15, 179 (1993); S. E. Smith, N. Durmuller, B. S. Meldrum, Eur. J. Pharmacol. 201, 179 (1991)]. AMPA receptor antagonists have also demonstrated promise in chronic neurodegenerative disorders such as Parkinsonism. [T. Klockgether, L. Turski, T. Honoré, Z. Zhang, D. M. Gash, R. Kurlan, J. T. Greenamyre, Ann. Neurol., 34(4), 585-593 (1993)].

Excitatory amino acid receptor antagonists that block NMDA receptors are also recognized for usefulness in the treatment of disorders. NMDA receptors are intimately involved in the phenomenon of excitotoxicity, which may be a critical determinant of outcome of several neurological disorders. Disorders known to be responsive to blockade of the NMDA receptor include acute cerebral ischemia (stroke or cerebral trauma, for example), muscular spasm, convulsive disorders, neuropathic pain and anxiety, and may be a

- 5 -

significant causal factor in chronic neurodegenerative disorders such as Parkinson's disease [T. Klockgether, L. Turski, Ann. Neurol. 34, 585-593 (1993)], human immunodeficiency virus (HIV) related neuronal injury, amyotrophic lateral sclerosis (ALS), Alzheimer's disease [P.T. Francis, N.R. Sims, A.W. Procter, D.M. Bowen, J. Neurochem. 60 (5), 1589-1604 (1993)] and Huntington's disease. [See S. Lipton, TINS 16 (12), 527-532 (1993); S.A. Lipton, P.A. Rosenberg, New Eng. J. Med. 330 (9), 613-622 (1994); and C.F. Bigge, Biochem. Pharmacol. 45, 1547-1561 (1993) and references cited therein.]. NMDA receptor antagonists may also be used to prevent tolerance to opiate analgesia or to help control withdrawal symptoms from addictive drugs (Eur. Pat. Appl. 488,959A).

An object of this invention is to provide novel substituted alicyclic ring fused 2,3-quinoxalinediones which function as either or both NMDA antagonists or non-NMDA antagonists.

A further object of this invention is to provide a pharmaceutical composition containing an effective amount of the substituted alicyclic fused ring 2,3-quinoxalinediones to treat cerebrovascular disorders responsive to blocking any or all of NMDA receptors, AMPA receptors and kainate-receptors.

Another object of this invention is to provide a method of treating disorders responsive to the antagonism of glutamate or aspartate receptors in a human by administering a pharmaceutically effective amount of the substituted alicyclic fused ring 2,3-quinoxalinediones of this invention.

- 6 -


Another object of this invention is to provide novel methods of preparing substituted alicyclic ring fused 2,3-quinoxalinediones.

- 5 A further object of this invention is directed to novel intermediates of the substituted alicyclic ring fused 2,3-quinoxalinediones of this invention.

SUMMARY OF THE INVENTION

10

The present invention is directed to compounds represented by the formula (I):



- 15 or a pharmaceutically acceptable salt thereof wherein

Z is an alicyclic fused ring having 5 to 7 carbon atoms;

R¹ is hydrogen, an alkyl or an arylalkyl;

- 20 X and Y are independently hydrogen, halogen, nitro, cyano, COOH, CONR²R³, SONR²R³ wherein R² and R³ are independently hydrogen, alkyl having 1 to 6 carbon atoms, cycloalkyl or aralkyl; and

- A is O, NR⁴, CH₂NR⁴, CN, tetrazole or CO wherein R⁴ is hydrogen, alkyl, alkanol, alkyl amine or aralkyl,
- 25 wherein

- (i) when A is O, NR⁴ or CH₂NR⁴ then B is hydrogen, alkyl, alkenyl, alkynyl, aryl, aralkyl, alkanol, alkoxy, alkyl amine, heterocyclic, alkylheterocyclic,
- 30 alkylcarbonyl, cycloalkylcarbonyl, arylcarbonyl, aralkylcarbonyl, heterocyclic-carbonyl, alkylheterocyclic-carbonyl, any of which may be unsubstituted or substituted by one or more hydroxy,

- 7 -

thiol, amine, alkyl or butoxycarbonyl group, CONR^5R^6 wherein R^5 is hydrogen, alkyl having 1 to 6 carbon atoms, aralkyl or NR^5 is a cyclic amine and R^6 is alkyl, aryl, aralkyl or not present when NR^5 is the cyclic
5 amine, or when A is NR^4 or CH_2NR^4 then B is a common amino acid moiety joined by an amide bond or B joins with R^4 to form a four to seven membered heterocyclic ring, provided that when Z is a fused cyclohexyl ring and R^4 is hydrogen then B is not hydrogen;

10 (ii) when A is CN then B is not present and Z is not a fused cyclohexyl ring;

(iii) when A is tetrazole then B is hydrogen or alkyl having 1 to 6 carbon atoms; and

(iv) when A is CO then B is hydroxy, alkoxy,
15 aralkoxy, alkyl having 1 to 6 carbon atoms, aralkyl, NR^7R^8 wherein R^7 is hydrogen, alkyl having 1 to 6 carbon atoms, aralkyl or NR^7 is a cyclic amine and R^8 is alkyl, aryl aralkyl or not present when NR^7 is the cyclic amine.

20

Examples of pharmaceutically acceptable addition salts include inorganic and organic acid addition salts such as the hydrochloride, hydrobromide, phosphate, sulphate, citrate, lactate, tartrate, maleate,
25 fumarate, mandelate, oxalate, and the acetate.

Alternatively, pharmaceutically acceptable inorganic and organic base addition salts may be used such as sodium hydroxide, potassium hydroxide, lithium hydroxide and the like.

30

Halogen is fluorine, chlorine, bromine, or iodine; fluorine, chlorine, and bromine are preferred groups.

Alkyl means a straight chained or branched chain of
35 from one to six carbon atoms or cyclic alkyl of from three to seven carbon atoms including, but not limited to methyl, ethyl, propyl, isopropyl, butyl, isobutyl,

- 8 -

t-butyl, pentyl, hexyl, cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

- Alkenyl means a straight chained or branched chain alkenyl group of two to six carbon atoms or a cyclic alkenyl group of three to seven carbon atoms, for example, but not limited to ethylene, 1,2- or 2,3-propylene, 1-2, 2,3-, or 3,4-butylene, cyclopentene, or cyclohexene.
- Alkynyl means a straight chained or branched chain alkynyl group of two to six carbon atoms, for example, but not limited to ethynyl, 2,3-propynyl, 2,3- or 3,4-butyne.
- Aryl means a monocyclic or bicyclic carbocyclic aromatic ring system, for example, but not limited to phenyl, 2-naphthyl, or 1-naphthyl.
- Aralkyl means aryl as defined above and alkyl as defined above, for example, but not limited to benzyl, 2-phenylethyl, 3-phenylpropyl; a preferred group is benzyl.
- Alkanol means alkyl as defined above substituted by a hydroxy group, for example, but not limited to methanol, ethanol, propanol, butanol, heptanol or hexanol.
- Alkoxy means an alkoxy group containing an alkyl group as defined above.
- Alkyl amine means alkyl as defined above substituted by an amino group, for example, but not limited to methyl amine, ethyl amine, propyl amine, butyl amine, heptyl amine or hexyl amine.

- 9 -

Heterocyclic means a monocyclic or bicyclic carbocyclic non-aromatic or aromatic 4 to 16 member ring system substituted by one or more hetero atoms, which can be the same or different, and includes, for example,
5 piperidinyl, piperazinyl, morpholinyl, pyrrolidinyl, thienyl, benzo[b]thienyl, naphtho[2,3[b]]thienyl, thianthrenyl, furyl, pyranyl, isobenzofuranyl, chromenyl, xanthenyl, phenoxanthiinyl, 2H-pyrrolyl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl,
10 pyrimidinyl, pyridazinyl, indoliziny, isoindolyl, 3H-indolyl, indolyl, indazolyl, purinyl, 4H-quinoliziny, isoquinolyl, quinolyl, phthalziny, naphthyridinyl, quinoxaliny, cinnolinyl, pteridinyl, 5aH-carbozoly, carbozoly, β -carbolinyl, phenanthridinyl, acridinyl,
15 perimidinyl, phenanthrolinyl, phenazinyl, isothiazolyl, phenothiazinyl, isoxazoly, furazanyl and phenoxazinyl groups. The heterocyclic group can be unsubstituted or substituted by, for example, hydroxy, thiol, amine, alkyl or butoxycarbonyl groups.

20

Common amino acid moiety means the naturally occurring α -amino acids.

The instant invention is also related to a
25 pharmaceutical composition containing the compound defined by formula I in an amount effective to treat cerebrovascular disorders responsive to the blockade of glutamate receptors, including either or both NMDA receptors and non-NMDA receptors (such as the
30 α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) receptor and the kainate receptor), and a pharmaceutically acceptable carrier. Exemplary disorders responsive to such treatment include cerebral ischemia caused by cerebral trauma, stroke,
35 hypoglycemia, heart attack, and surgery; anxiety and schizophrenia; and chronic neurodegenerative disorders such as Huntington's disease, ALS, Parkinsonism and


- 10 -

Alzheimer's disease. The pharmaceutical composition of this invention may also be employed as an analgesic or the treatment of epilepsy.


- 5 The invention further relates to a method of treating cerebrovascular disorders responsive to antagonism of glutamate receptors including either or both NMDA receptors and non-NMDA receptors by administering a compound of above-defined formula I in a unit dosage
10 form.

The invention is also related to a method for preparing the compound of formula (I) comprising the steps of:

- 15 (a) reacting a compound of formula



- wherein Z is an alicyclic fused ring having 5 to 7 carbon atoms and A' is OH, N₃, COOH, COO-alkyl, COO-benzyl, a heterocyclic ring or a protected amine, with
20 a halogenating reagent to produce a compound of formula




wherein X is a halogen,


25

- (b) nitrating the compound formed in step (a) to produce a compound of formula

- 11 -




- (c) sequentially hydrolyzing an acetamide of the compound of step (b) and then catalytically
 5 hydrogenating the hydrolyzed compound to produce a compound of formula




- wherein X' is a halogen or hydrogen,
 10

- (d) condensation of the compound of step (c) with oxalic acid or an oxalic acid ester to produce a compound of formula




15

- (e) optionally hydrogenating the compound of step (d) when X' is halogen to produce a compound of formula




- 20 (f) optionally reacting the compound of steps (d) or (e) with an electrophilic substituent to produce a compound of formula

- 12 -



wherein X and Y are independently hydrogen, halogen, nitro, cyano, COOH, CONR²R³, SONR²R³, wherein R² and R³ are independently hydrogen, alkyl, cycloalkyl or aralkyl; and

(g) optionally reacting the compound of steps (d), (e) or (f) when A' is amine or OH with a reagent selected from the group consisting of electrophilic reagents and hydride reagents to produce a compound of formula



wherein A and B are as defined for formula (I).

15

Typically the halogenating agent employed in the methods of this invention is selected from the group consisting of bromine, N-bromosuccinimide, chlorine, and iodochloride or the like. When a brominating agent is employed it is preferably present in an acetic acid/sodium acetate solution. The halogenation step is generally carried out at a temperature of about 15-50°C for about 15 minutes to 24 hours.

25 Either fuming nitric acid or potassium nitrate in an acid selected from the group consisting of trifluoroacetic acid, acetic acid, concentrated sulfuric acid or the like are generally employed during the steps of nitrating in the methods described herein.

- 13 -

More particularly, fuming nitric acid in trifluoroacetic acid, acetic acid or the like is employed at about 0°C to room temperature for about 15 minutes to about 8 hours. Alternatively, fuming nitric acid, potassium nitrate or the like in concentrated sulfuric acid may be used.

Hydrogenation in the methods of this invention is generally accomplished with a nickel or palladium catalyst in a hydrogen atmosphere. For example, deactivated Raney nickel can be used in a hydrogen atmosphere of about 1 atm to 100 psi in a solvent such as methanol, ethanol or tetrahydrofuran or the like. Such a reaction is normally carried out at ambient temperature to about 100°C. It is also possible to employ a more powerful catalyst, such as 20% palladium on carbon under similar conditions in order to dehalogenate the intermediates of this invention.

The step of condensation in the methods of this invention is generally accomplished with oxalic acid or an oxalic acid ester such as dimethyl oxalate or the like. The condensation reaction is preferably carried out at a temperature of about 25-150°C for about 30 minutes to 24 hours in an aqueous acidic media or an organic solvent such as acetonitrile, methanol or the like.

The electrophilic reagents typically employed in the methods of this invention are selected from the group consisting of alkyl halides, acylhalide, aldehydes, isocyanates and mixtures thereof. These reagents are generally used in a solvent such as methanol, dimethylformamide or the like at a temperature of about 0 to 100°C. A hydride reagent such as sodium cyanoborohydride or the like may be employed in the methods of this invention when a reductive step is

- 14 -


desired such as with reductive amination of an intermediate.

5 The invention is further related to a method for preparing substituted cyclopentyl fused ring quinoxalinediones comprising the steps of:


(a) acetylating and nitrating a compound of the formula



to form a compound of the formula




15 (b) hydrogenation of a nitro group, acetylating a resulting aniline and halogenating the compound formed in step (a) to form a compound of the formula



20 wherein X is a halogen,


(c) nitrating the compound of step (b) to form a compound of the formula




25

(d) hydrolysis of the compound of step (c) to form a compound of the formula

- 15 -



(e) hydrogenating the compound of step (d) to form a diamine compound of the formula




5

or


reacting the compound of step (d) with XCCOOR^{16} , $\begin{array}{c} \text{O} \\ || \end{array}$

wherein R^{16} is hydrogen, alkyl or aralkyl, to form an amide compound of the formula



10

(f) optionally hydrogenating the diamino compound of step (d) to form a compound of the formula




15

(g) condensation of the diamino compound of step (d) or the compound of step (f) with oxalic acid or an


- 16 -

oxalic acid ester or hydrogenation of the amide compound of step (e) to form a compound of the formula



5 wherein X' is a halogen or hydrogen, or


(h) optionally reacting the compound of step (g) with an electrophilic substituent to produce a compound of formula



10

wherein X and Y are the same as defined for formula (I), and

15 (i) optionally hydrolyzing compound of steps (g) or (i) with a reagent selected from the group consisting of electrophilic reagents and hydride reagents to produce a compound of the formula



20



wherein R is hydrogen, alkyl, alkenyl, alkynyl, aryl, aralkyl, alkylheterocyclic, alkylcarbonyl, aralkylcarbonyl, alkylheterocycliccarbonyl, CONR⁵R⁶

- 17 -

wherein R^5 is hydrogen, alkyl, aralkyl or NR^5 is a cyclic amine and R^6 is alkyl, aryl or aralkyl, or a common amino acid moiety joined by an amide bond.


- 5 Generally the step of acetylating in this method is performed with acetyl chloride or acetic anhydride at a temperature of about 25 to 80°C for about 5 minutes to 2 hours. Hydrolysis of the acetamide intermediate in this process can be accomplished with 2N HCL or the
 10 like. Other steps in the methods described herein may be readily accomplished by those skilled in the art.

This invention is further directed to novel intermediates which may be prepared during the
 15 preparation of the substituted alicyclic ring fused quinoxalinediones of this invention. The novel intermediate compounds are represented by the formulae II, III, IV and V:



- wherein R^9 is N_3 or $R^{10}HN$ — wherein R^{10} is hydrogen alkyl, alkenyl, alkynyl, aryl, aralkyl, alkylheterocyclic,
 25 alkylcarbonyl, aralkylcarbonyl, alkylheterocycliccarbonyl, $CONR^5R^6$ wherein R^5 is hydrogen, alkyl, aralkyl or NR^5 is a cyclic amine and R^6 is alkyl, aralkyl, or a common amino acid moiety joined by an amide bond,

- 18 -




(IV)

wherein A' is OH, N, or R¹⁰NH— and R¹⁰ is as previously described, Z is an alicyclic fused ring having 5 to 7 carbon atoms, R¹¹ and R¹² are independently NHAc,

NHCCOOR^{17} , NH₂ or NO₂ and R¹³ and R¹⁴ are independently

hydrogen or Br provided that at least one of R¹³ and R¹⁴ is Br, R¹⁷ is hydrogen, alkyl or aryl, and



(V)

10

wherein A' and Z are as previously described provided that when Z is a fused cyclohexyl ring R¹⁰ is not hydrogen, and R¹, X and wherein Y are as described for formula (I). Preferred intermediates of this invention illustrated by formula (IV) include those where when Z is a fused cyclohexyl ring, R¹¹ is NHAc or NH₂, R¹² is NO₂ or NH₂, R¹³ is hydrogen and R¹⁴ is Br and when Z is a fused cyclopentyl ring R¹¹ is NO₂ or NH₂, R¹² is NHAc or NH₂, R¹³ is Br and R¹⁴ is hydrogen. Preferred intermediates illustrated by formula (V) include those where Z is a fused cyclopentyl or cyclohexyl ring.

- 19 -

DETAILED DESCRIPTION OF
THE PREFERRED EMBODIMENTS


The substituted alicyclic ring fused quinoxalinediones
5 of this invention are represented by previously defined
formula I.

Preferably, X and Y are independently hydrogen, bromo
and nitro. It is also preferred that R¹ is a hydrogen
10 and Z is a fused cyclopentyl or cyclohexyl ring, either
of which are substituted by hydroxy, benzyloxy,
acetamide, benzyl amino, piperidine-4-carboxylic acid
tert-butylester, piperidine-1-carboxylic acid amide,
benzamide and cyclohexylamide groups.


15

More particularly, when A is O then preferably B is
hydrogen or a benzyl. When A is NR⁴ or CH₂NR⁴, then
preferably R⁴ is hydrogen, methyl or ethyl, and B is
methyl, ethyl, acetyl,

20



- 20 -



- 5 It is also preferred that when A is NR^4 or CH_2NR^4 then B may join with R^4 to form a heterocyclic ring, more preferably a pyrrolidine or piperidine ring.

10 Exemplary preferred compounds of Formula I include, without limitation:

9-benzyloxy-6-bromo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

- 15 6-bromo-9-hydroxy-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

20 piperidine-4-carboxylic acid (6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-octahydro-benzo[f]quinoxalin-9-yl) amide;

N-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-octahydro-benzo[f]quinoxalin-9-yl)-acetamide;

- 21 -

9-benzylamino-6-bromo-1,4,7,8,9,10-hexahydro-
benzo[f]quinoxaline-2,3-dione;

5 N-(5-bromo-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-
cyclopenta[f]-quinoxalin-8-yl)-acetamide;

N-(5-bromo-6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-
cyclopenta[f]quinoxalin-8-yl)-acetamide;

10 N-(2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-
cyclopenta[f]quinoxalin-8-yl)-acetamide;

N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-
cyclopenta[f]quinoxalin-8-yl)-acetamide;

15 8-amino-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-
2,3-dione hydrochloride;

20 4-(2,3,-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]-
quinoxalin-8-ylcarbamoyl)-piperidine-1-carboxylic acid
tert-butyl ester;

25 4-(6-nitro-2,3,-dioxo-2,3,4,7,8,9-hexahydro-1H
cyclopenta[f]quinoxalin-8-yl-carbamoyl)-piperidine
hydrochloride;

N-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-
octahydrobenzo[f]quinoxalin-9-yl)-benzamide;

30 N-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-
octahydrobenzo[f]quinoxalin-9-yl) cyclohexylamide;

35 8-amino-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]-
quinoxaline-2,3-dione hydrochloride;

N-(6-nitro-2,3-dioxo-4,7,8,9-tetrahydro-1H
cyclopenta[f]quinoxalin-8-yl)-benzamide;

- 22 -

4- (6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-
octahydrobenzo[f]quinoxalin-9-ylcarbamoyl)-piperidine
hydrochloride;

5 2- (6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-
octahydrobenzo[f]quinoxalin-9-ylcarbamoyl) ethyl- (4-
hydroxy) benzene;

10 N-phenyl-N' - (6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10 -
octahydrobenzo[f]quinoxalin-9-yl)-urea;

N-methyl-N- (6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-
octahydro-benzo[f]quinoxalin-9-yl)-benzamide;

15 thiophene-2-carboxylic acid (6-nitro-2,3-dioxo-
2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-
yl)-amide;

20 furan-2-carboxylic acid (6-nitro-2,3-dioxo-2,3,4,7,8,9-
hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-amide;

thiophene-3-carboxylic acid (6-nitro-2,3-dioxo-
2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-
yl)amide;

25 N- (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro
-1H-cyclopenta[f]quinoxalin-8-yl)-nicotinamide;

30 pyridine-2-carboxylic acid (6-nitro-2,3-dioxo-
2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-
amide;

35 2,3,4,5-tetrahydroxy-tetrahydro-furan-2-carboxylic acid
(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-
cyclopenta[f]quinoxalin-8-yl)-amide;

- 23 -

- pyrrolidine-2-carboxylic acid (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-amide;
- 5 benzo[b]thiophene-2-carboxylic acid (6-nitro-2,3,-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-amide;
- 10 2-(1H-indol-3-yl)-N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide;
- N-methyl thiophene-2-carboxylic acid-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-amide;
- 15 N-methyl-N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-benzamide;
- 20 6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-carboxylic acid methyl-phenyl-amide;
- 25 6-nitro-2,3,-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-carboxylic acid;
- 6-nitro-2,3,-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-carboxylic acid phenylamide;
- 30 9-methylamino-6-nitro-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;
- 6-nitro-9-pyrrolidin-1-yl-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;
- 35 8-dimethylamino-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

- 24 -

8-methylamino-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione methane sulfonate salt;

5 2-[bis-(2-hydroxy-ethyl)-amino]-N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide;

10 6-nitro-8-pyrrolidin-1-yl-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

N-methyl-N-(6-nitro-2,3,-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-yl)-acetamide;

15 6-nitro-8-piperidin-1-yl-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

8-diethylamino-6-nitro-4,7,8,9,-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

20 6-nitro-8-(thiophen-2-ylmethoxy)-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

25 6-nitro-9-(thiophen-2-ylmethoxy)-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

30 6-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yloxymethyl)-1,2,3,4-tetrahydro-isoquinoline-3-carboxylic acid;

6-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yloxymethyl)-1,2,3,4-tetrahydro-isoquinoline-3-carboxylic acid;

35 6-nitro-8-[2-(1H-pyrrol-2-yl)-ethoxy]-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

- 25 -

6-nitro-9-[2-(1H-pyrrol-2-yl)-ethoxy]-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

5 6-nitro-8-([(1H-tetrazol-5-ylmethyl)-amino]-methyl)-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

10 6-nitro-9-([(1H-tetrazol-5-ylmethyl)-amino]-methyl)-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

6-nitro-8-[(1H-tetrazol-5-ylmethyl)-amino]-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

15 6-nitro-8-[(1H-tetrazol-5-ylmethyl)-amino]-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

20 8-{[methyl-(1H-tetrazol-5-ylmethyl)-amino]-methyl}-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

25 9-{[methyl-(1H-tetrazol-5-ylmethyl)-amino]-methyl}-6-nitro-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

6-nitro-8-[2-(1H-tetrazol-5-yl)-ethoxy]-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

30 6-nitro-9-[2-(1H-tetrazol-5-yl)-ethoxy]-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

35 6-nitro-8-[2-(1H-tetrazol-5-yl)-ethylamino]-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

6-nitro-9-[2-(1H-tetrazol-5-yl)-ethylamino]-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

- 26 -

6-nitro-8-(1H-tetrazol-5-ylmethoxy)-4,7,8,9-tetrahydro-
1H-cyclopenta[f]quinoxaline-2,3-dione;

5 6-nitro-9-(1H-tetrazol-5-ylmethoxy)-1,4,7,8,9,10-
hexahydro-benzo[f]quinoxaline-2,3-dione;

10 N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-
cyclopenta[f]quinoxalin-8-ylmethyl)-2-(1H-tetrazol-5-
yl)-acetamide;

N-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-
benzo[f]quinoxalin-9-ylmethyl)-2-(1H-tetrazol-5-yl)-
acetamide;

15 2-(3-hydroxy-5-methyl-isoxazol-4-yl)-N-(6-nitro-2,3-
dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-
8-yl)-acetamide;

20 2-(3-hydroxy-5-methyl-isoxazol-4-yl)-N-(6-nitro-2,3-
dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl)-
acetamide;

25 2-(3-hydroxy-5-methyl-isoxazol-4-yl)-N-(6-nitro-2,3-
dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-
8-ylmethyl)-acetamide;

30 2-(3-hydroxy-5-methyl-isoxazol-4-yl)-N-(6-nitro-2,3-
dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl
methyl)-acetamide;

8-(3-hydroxy-5-methyl-isoxazol-4-ylmethoxy)-6-nitro-
4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-
dione;

35 9-(3-hydroxy-5-methyl-isoxazol-4-ylmethoxy)-6-nitro-
1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

- 27 -

- N- (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl) -2- (1H-tetrazol-5-yl) -acetamide;
- 5 N- (6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl) -2- (1H-tetrazol-5-yl) -acetamide;
- 10 3- [(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-ylmethyl) -carbamoyl] -1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid;
- 15 3- [(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-ylmethyl) -carbamoyl] -1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid;
- 20 8- [2- (3-hydroxy-5-methyl-isoxazol-4-yl) -ethoxy] -6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;
- 25 9- [2- (3-hydroxy-5-methyl-isoxazol-4-yl) -ethoxy] -6-nitro-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;
- 2- (2-hydroxy-phenyl) -N- (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl) -acetamide;
- 30 2- (2-hydroxy-phenyl) -N- (6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl) -acetamide;
- 2-hydroxy-N- (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-ylmethyl) -benzamide;
- 35 2-hydroxy-N- (6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-ylmethyl) -benzamide;

- 28 -

6-nitro-8-(2-pyridin-2-yl-ethoxy)-4,7,8,9-tetrahydro-
1H-cyclopenta[f]quinoxaline-2,3-dione;

5 6-nitro-9-(2-pyridin-2-yl-ethoxy)-1,4,7,8,9,10-
hexahydro-benzo[f]quinoxaline-2,3-dione;

10 N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-
cyclopenta[f]quinoxalin-8-yl)-2-[2-(1H-tetrazol-5-yl)-
phenyl]-acetamide;

N-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-
benzo[f]quinoxalin-9-yl)-2-[2-(1H-tetrazol-5-yl)-
phenyl]-acetamide;

15 1-methyl-2,4-dioxo-1,2,3,4-tetrahydro-pyrimidine-5-
carboxylic acid (6-nitro-2,3-dioxo-2,3,4,7,8,9-
hexahydro-1H-cyclopenta[f]quinoxalin-8-ylmethyl)-amide;

20 1-methyl-2,4-dioxo-1,2,3,4-tetrahydro-pyrimidine-5-
carboxylic acid (6-nitro-2,3-dioxo-1,4,7,8,9,10-
hexahydro-benzo[f]quinoxalin-9-ylmethyl)-amide;

25 6-nitro-8-(5-oxo-4,5-dihydro-[1,3,4]oxadiazol-2-
ylmethoxy)-4,7,8,9-tetrahydro-1H-
cyclopenta[f]quinoxaline-2,3-dione;

6-nitro-9-(5-oxo-4,5-dihydro-[1,3,4]oxadiazol-2-
ylmethoxy)-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-
2,3-dione;

30 6-nitro-8-(pyridin-2-ylamino)-4,7,8,9-tetrahydro-1H-
cyclopenta[f]quinoxaline-2,3-dione;

35 6-nitro-9-(pyridin-2-ylamino)-1,4,7,8,9,10-hexahydro-
benzo[f]quinoxaline-2,3-dione;

- 29 -

- 4 - {[(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-ylmethyl)-amino]-methyl}-benzoic acid;
- 5 4 - {[(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-ylmethyl)-amino]-methyl}-benzoic acid;
- 10 2-hydroxy-benzoic acid 6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl ester;
- 2-hydroxy-benzoic acid 6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl ester;
- 15 N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-phthalamic acid;
- N-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl)-phthalamic acid;
- 20 2-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yloxy)-benzoic acid;
- 25 2-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl oxy)-benzoic acid;
- 30 (3-hydroxy-5-methyl-isoxazol-4-yl)-acetic acid 6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl ester;
- (3-hydroxy-5-methyl-isoxazol-4-yl)-acetic acid 6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl ester;
- 35 2-hydroxy-N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-benzamide; and


- 30 -

2-hydroxy-N-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydrobenzo[f]quinoxalin-9-yl)-benzamide.

This invention is also directed to methods for
 5 preparing the substituted alicyclic fused ring
 2,3-quinoxalinedione compounds of formula I. One
 embodiment of this invention is directed to the method
 employing the exemplary reaction schemes set forth in
 Schemes II and III employing starting materials
 10 prepared, for example, via the method set forth in
 Scheme I. Yet another embodiment of this invention is
 directed to the method of preparing the substituted
 cyclopentyl fused ring compounds of formula I via the
 reaction Scheme IV. The starting materials employed in
 15 these reaction schemes are either readily available,
 can be prepared by known methods or by the method set
 forth in Scheme I.

Scheme I

20



The tetrahydronaphthalene derivatives substituted on
 the saturated ring, as shown in Scheme 1, are readily
 25 available or prepared by those skilled in the art. For
 example, 1-amino-7-hydroxynaphthalene can be
 hydrogenated at a pressure from about 50 to about 1000
 psi and at a temperature of about 20°C to about 100°C
 to form the tetrahydronaphthalene adduct, which can
 30 then act as a starting material for hydroxy and ether
 adducts, or via conversion to a leaving group such as a

- 31 -

mesylate, tosylate, halide or the like, can be further elaborated to an azide by treatment with sodium azide, or displaced by an amine by nucleophilic substitution. Stereochemistry at the chiral center can be controlled
5 by means known to those skilled in the art, such as Mitsunobu inversion, or by separation via chiral salts or chiral chromatography, or by reacting with a chiral protecting group and separating the resulting disastereomeric mixture.


10

Alternatively, a β -tetralone (or α -tetralone) derivative can be reduced directly to the alcohol using any number of reducing agents such as sodium borohydride, or converted to the carboxylic acid adduct
15 by formation of the cyanohydrin, followed by elimination and hydrolysis. Subsequent treatment with diphenylphosphoryl azide or the like (Schmidt rearrangement), followed by heating and alcoholysis (or hydrolysis) can lead to either the carbamate or
20 unprotected amine.

Another embodiment of this invention is directed to a second novel process of preparing the compound of Formula I in the following manner.

25

Scheme II set forth below illustrates a method of preparing novel intermediates and the substituted alicyclic fused ring 2,3-quinoxalinediones of this invention employing starting materials that are either
30 readily available or prepared via Scheme I.


Scheme II

As shown in Scheme II wherein A' is OH, N₃, a
 5 heterocyclic ring or a protected amine, the aromatic
 ring can be manipulated by sequential electrophilic
 aromatic substitution reactions, bromination and
 nitration, to obtain the desired substitution pattern.
 Reduction of the nitro functionality via catalytic
 10 hydrogenation can be done under conditions that give
 the diamine with the bromine intact, or give the
 diamine and dehalogenation. Subsequent condensation
 with oxalic acid provides the benzo(f)quinoxalinedione.
 Hydrogenolysis of the bromine followed by electrophilic
 15 aromatic substitution provides aryl substituted
 derivatives.

- 33 -


The compounds of this invention may be prepared employing the compound prepared via Scheme II. This is illustrated by Scheme III below.

5 Scheme III



When A' is OH or NH₂, the alcohol or amine can be reacted with an electrophilic reagent, such as an alkyl halide, mesylate or tosylate or the like, or an acid chloride, acylimidazole or the like, to generate
 10 ethers, esters, secondary amine or amides. When A' is COOH, it can be activated as an acylimidazole, or the like, with carbonyldiimidazole (or dicyclohexylcarbodiimide) and then reacted with amines
 15 or alcohols to form the reverse amide or ester. When X,Y are nonreactive species, hydride reagents can be used to reduce the amide derivatives to amines; in the case where A' is NHCO- the amine is directly attached to the cyclohexyl ring, and when A' is CONR-, a
 20 methylene spacer separates the amine from the cyclohexyl ring. Preparation of analogs that incorporate a fused cyclopentyl ring rather than the cyclohexyl ring can be accomplished in a similar manner.

25 Yet another embodiment of this invention is directed to the process of preparing substituted cyclopentyl fused ring 2,3-quinoxalinediones compounds of Formula I as shown below.

Scheme IV


The disposition of the functional groups on the aromatic ring in the cyclopentyl ring fused analogs can be established by traditional aromatic electrophilic substitution reactions as shown in Scheme IV.

Introduction of the nitro functionality followed by reduction and acetylation of the aniline allows bromination to proceed primarily in the least hindered positions ortho to the acetamide. Nitration is then directed to the open ortho position which establishes the correct regiochemistry for the entire aromatic ring. After formation of the quinoxalinedione, the final substitution pattern on the aromatic ring can be established by standard chemistry known to those skilled in the art. The exo amino functionality on the cyclopentyl ring can be manipulated as shown before in Scheme III.


A preferred method for preparing the compound of formula (I) comprises the steps of:

- 35 -


(a) reacting a compound of formula



wherein Z is an alicyclic fused ring having 5 to 7
5 carbon atoms and A' is OH, N₃, COOH, COO-alkyl,
COO-benzyl, a heterocyclic ring such as a pyrrolidine
or a piperidine, or a protected amine such as
phthalimide, acetamide or a carbamate (e.g., Boc or
Cbz), with a brominating agent to produce a compound of
10 formula




(b) nitrating the compound formed in step (a) to
produce a compound of formula



15


(c) catalytically hydrogenating the compound of step
(b) to produce a compound of formula



20


- 36 -

(d) condensation of the compound of step (c) with oxalic acid or an oxalic acid ester such as dimethyl oxalate or the like, to produce a compound of formula




5

(e) optionally hydrogenating the compound of step (d) to produce a compound of formula




10 (f) optionally reacting the compound of steps (d) or (e) with an electrophilic substituent to produce a compound of formula



15 wherein X and Y are independently hydrogen, halogen, nitro, cyano, COOH, CONR²R³, SONR²R³, wherein R² and R³ are independently hydrogen, alkyl, cycloalkyl or aralkyl; and

- 37 -

(g) optionally reacting the compound of steps (d), (e) or (f) wherein A' is NR⁴H or OH with a reagent selected from the group consisting of electrophilic reagents and hydride reagents, to produce a compound of formula



5

wherein A, B and R⁴ are as defined for formula (1). Exemplary electrophilic reagents include alkyl halides, mesylates, tosylates, acid chlorides, acylimidazoles, isocyanates, aldehydes and the like. When A' is COOH, it can be activated with a coupling reagent such as CDI, DCC, thionyl chloride and the like and further reacted with an amine, including natural amino acids, or alcohols, to obtain the corresponding amides and esters. In this preferred method A' may also be a piperidine or pyrrolidine ring.


10

15


The preparation of substituted cyclopentyl fused ring quinoxalinediones can be preferably accomplished by:

20

(a) acetylating and nitrating a compound of the formula




25 to form a compound of the formula




- 38 -


(b) reducing, acetylating and brominating the compound formed in step (a) to form a compound of the formula




5 (c) nitrating the compound of step (b) to form a compound of the formula



10 (d) hydrogenating the compound of step (c) to form a compound of the formula




(e) optionally hydrogenating the compound of step (d) to form a compound of the formula



15


(f) condensation of the compound of steps (d) or (e) with oxalic acid to form a compound of the formula



20

- 39 -


(g) optionally reacting the compound of step (f) with an electrophilic substituent to produce a compound of formula



5

wherein X and Y are the same as defined for formula (I), and

(h) optionally reacting the compound of steps (f) or (g) with a reagent selected from the group consisting of electrophilic reagents and hydride reagents, to produce a compound of the formula



wherein R is hydrogen, alkyl, alkenyl, alkynyl, aryl, aralkyl, alkylheterocyclic, alkylcarbonyl, aralkylcarbonyl, alkylheterocycliccarbonyl, CONR⁵R⁶ wherein R⁵ is hydrogen, alkyl, aralkyl or NR⁵ is a cyclic amine and R⁶ is alkyl, aryl, aralkyl or not present, or a common amino acid moiety joined by an amide bond.

The methods set forth herein, and particularly as illustrated in Schemes I, II, III and IV, may also be employed to prepare novel intermediates of this invention. The preferred novel intermediates include:

- 40 -

acetic acid 8-acetylamino-5-bromo-7-nitro-1,2,3,4-tetrahydronaphthalen-2-yl ester;

5 8-amino-5-bromo-7-nitro-1,2,3,4-tetrahydronaphthalen-2-ol;

7,8-diamino-5-bromo-1,2,3,4-tetrahydronaphthalen-2-ol;

10 methanesulfonic acid 8-acetyl-1,2,3,4-tetrahydronaphthalen-2-yl ester;

N-(7-azido-5,6,7,8-tetrahydronaphthalene-1-yl)-acetamide;

15 N-(7-azido-4-bromo-5,6,7,8-tetrahydronaphthalen-1-yl)-acetamide;

20 N-(7-azido-4-bromo-2-nitro-5,6,7,8-tetrahydronaphthalen-1-yl) acetamide;

7-azido-4-bromo-2-5,6,7,8-tetrahydro-naphthalen-1-ylamine;

25 N-7-benzyl-4-bromo-2-nitro-5,6,7,8-tetrahydronaphthalene-1,7-diamine;

N-(5-acetylamino-6-bromo-indan-2-yl)-acetamide;

30 N-(5-acetylamino-6-bromo-4-nitro-indan-2-yl)-acetamide;

N-(5-amino-6-bromo-4-nitro-indan-2-yl)-acetamide;

6-bromo-4-nitro-indan-2,5-diamine monohydrochloride;
and

35

N-(4,5-diamino-6-bromo-indan-2-yl)-acetamide.

- 41 -

The compounds of the invention exhibit valuable biological properties because of their strong excitatory amino acid (EAA) antagonizing properties at one of several binding sites on glutamate receptors:

5 the AMPA ((RS)--amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (or kainic acid) binding site on AMPA (non-NMDA) receptors or the glycine site of NMDA receptors. The compounds generally have activity at both NMDA and non-NMDA receptors, and thus may act

10 in disorders arising from over excitation of either receptor family.

The compounds of the present invention exhibit binding affinity for the AMPA receptors as described by

15 Honoré T., et al., Neuroscience Letters 1985; 54:27-32. Preferred compounds demonstrate IC_{50} values $< 100 \mu M$ in this assay. The compounds of the present invention exhibit binding affinity for the kainate site (non-NMDA receptor) as described by London, ED and Coyle, J, Mol.

20 Pharmacol., 1979; 15:492. The compounds of the present invention exhibit binding affinity for the glycine site of the NMDA receptor as described by Jones, SM et al., Pharmacol. Methods 1989; 21:161. To functionally measure AMPA antagonist activity, the effects of the

25 agent on AMPA-induced neuronal damage in primary cortical neuronal cultures was examined using techniques similar to those outlined by Koh, J.-Y. et al., J. Neurosci., 1990; 10:693. The neuronal damage produced by long-term exposure to $100 \mu M$ AMPA was

30 measured by the release of the cytosolic enzyme lactate dehydrogenase (LDH).

Selected compounds of the present invention were tested by one or more of the above-described assays. The data

35 obtained in these assays is set forth in Tables 1 and 2. The IC_{50} values set forth in Tables 1 and 2 are a measure of the concentration (μM) of the test

- 42 -

substance which inhibits 50% of an induced release from the tested receptors.

Table 1. Cyclohexyl-fused Analogs

5

10

15

20

| Compound of Example | AMPA (IC ₅₀ μM) | Kainate (IC ₅₀ μM) | Glycine (IC ₅₀ μM) | LDH (IC ₅₀ μM) |
|------------------------|-------------------------------|----------------------------------|----------------------------------|------------------------------|
| (7) | 17 | 3 | 1.5 | |
| (8) | 14 | >100 | | |
| (15) | 0.7 | 2.8 | 1.8 | 20 |
| (17) | 0.56 | 2.2 | 0.62 | >30 |
| (19) | 25 | >100 | >100 | |
| (36) | 0.42 | 0.9 | 0.09 | 30 |
| (37) | 1.1 | 1.4 | 0.5 | 28 |
| (40) | 0.27 | 0.82 | 0.28 | 25 |
| (41) | 0.96 | 1.4 | >1 | |
| (42) | 0.15 | 0.83 | 0.18 | |
| (52) | 0.79 | 2.8 | >1 | |
| (53) | 1.1 | 1.6 | | |
| (86) | 2.2 | 3.8 | | |

Table 2. Cyclopentyl-fused Analogs

| Compound of Example | AMPA (IC ₅₀ μM) | Kainate (IC ₅₀ μM) | Glycine (IC ₅₀ μM) | LDH (IC ₅₀ μM) |
|------------------------|-------------------------------|----------------------------------|----------------------------------|------------------------------|
| (28) | >100 | >100 | | |
| (29) | 5.1 | | | |
| (30) | 25 | 87 | | |
| (32) | 0.4 | 0.9 | 1.3 | |
| (35) | 0.12 | 0.28 | <1 | |
| (38) | 2.6 | 2.1 | | |
| (39) | 0.11 | 0.48 | 0.4 | 18 |
| (54) | 0.05 | 0.18 | 0.3 | |
| (59) | 0.14 | 0.38 | >1 | 14 |
| (60) | 0.10 | 0.36 | 0.4 | 14 |
| (61) | 0.20 | 1.03 | >1 | 28 |
| (62) | 0.56 | 1.9 | >1 | |
| (63) | 0.58 | 1.1 | >1 | >30 |
| (64) | 0.64 | 0.86 | >1 | |
| (65) | 1.3 | 8.1 | | |
| (66) | 1.3 | 2.3 | | |
| (67) | 3.6 | 1.6 | | |
| (68) | 7.4 | 1.9 | | |
| (69) | 0.42 | 1.6 | >100 | |
| (78) | 0.42 | 1.9 | 14 | |
| (80) | 1.6 | 3.9 | | |
| (90) | 0.91 | 2.6 | 12 | |
| (100) | 1.2 | 2.2 | | |
| (101) | 1.1 | 1.4 | | |
| (104) | 1.5 | 2.2 | | |
| (105) | 2.2 | 2.2 | | |
| (108) | 3.4 | 1.6 | | |
| (109) | 5.1 | 5.4 | | |

35

Additionally, as a preliminary indicator of in vivo CNS activity related to anticonvulsant activity and potential neuroprotection, a maximal electroshock assay

- 44 -

in CF-1 strain mice (20-25 g) was performed with corneal electrodes by conventional methods as described previously (Krall et al., Epilepsia 1988; 19:409-428. The compounds of this invention generally demonstrated
5 ED₅₀ values of < 50 mg/kg.

The compounds of the invention, together with a conventional adjuvant, carrier, or diluent, may be placed into the form of pharmaceutical compositions and
10 unit dosages thereof, and in such form may be employed as solids, such as tablets or filled capsules, or liquids such as solutions, suspensions, emulsions, elixirs, or capsules filled with the same, all for oral use, in the form of suppositories for rectal
15 administration; or in the form of sterile injectable solutions for parenteral (including subcutaneous) use. Such pharmaceutical compositions and unit dosage forms thereof may comprises conventional ingredients in conventional proportions, with or without additional
20 active compounds or principles, and such unit dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily dosage range to be employed. Tablets containing 10 mg of active ingredients or, more broadly, 0.1 to 100 mg
25 per tablet, and accordingly suitable representative unit dosage forms.

Solid forms of pharmaceutical compositions for PO administration and injectable solutions are preferred.
30

The compounds of this invention are extremely useful in the treatment of central nervous system disorders related to their biological activity. The compounds of this invention may accordingly be administered to a
35 subject, including a human, in need of treatment, alleviation, or elimination of an indication associated with the biological activity of the compounds. This

- 45 -


includes especially excitatory amino-acid-dependent psychosis, excitatory amino-acid-dependent anoxia, excitatory amino-acid-dependent ischemia, excitatory amino-acid-dependent Parkinsonism, excitatory amino-acid-dependent convulsions, and excitatory amino-acid-dependent migraine. Suitable dosage ranges are 0.1 to 1000 mg daily, 10 to 50 mg daily, and especially 30 to 100 mg daily, dependent as usual upon the exact mode of administration, form in which administered, the indication toward which the administration is directed, the subject involved, and the body weight of the subject involved, and further, the preference and experience of the physician or veterinarian in charge.

The examples which follow are intended as an illustration of certain preferred embodiments of the invention, and no limitation of the invention is implied.

20

Example 1

Preparation of N-(7-hydroxy-5,6,7,8-tetrahydronaphthalen-1-yl)-acetamide




25

A mixture of 1-amino-7-hydroxynaphthalene (16.5 g, 0.1 mol), lithium hydroxide hydrate (4.3 g) and 20% palladium on carbon (4 g) was suspended in 600 mL 4:1 methanol and water and shaken on a Parr hydrogenation apparatus under a hydrogen atmosphere (52 psi) for about one hour. The hydrogen atmosphere was then recharged, and the mixture shaken an additional hour. After filtration through celite, the filtrate was concentrated and the pH adjusted with aqueous HCl to 7. Acetic anhydride (15 g) was added and swirled for 15

min. The solution was saturated with sodium chloride and extracted with ethyl acetate (3 x). The combined organic layer was dried over magnesium sulfate, filtered and evaporated to give the product (9.5 g) in 46% yield.

Example 2


Preparation of acetic acid 8-acetylamino-1,2,3,4-tetrahydronaphthalen-2-yl ester



A mixture of N-(7-hydroxy-5,6,7,8-tetrahydronaphthalen-1-yl)-acetamide (3 g, 15 mmol), acetic anhydride (2 g, 200 mmol) in pyridine (50 mL) was stirred at room temperature for 18 h. Solvent was removed by rotoevaporation and the residue was purified by column chromatography on silica gel (1:1 ethyl acetate:hexane as eluant) to give the product (2.2 g) as a tan solid in 59% yield.

Example 3

Preparation of acetic acid 8-acetylamino-5-bromo-1,2,3,4-tetrahydronaphthalen-2-yl ester




A mixture of acetic acid 8-acetylamino-1,2,3,4-tetrahydronaphthalen-2-yl ester (2.2 g, 8.9 mmol) and bromine (1.7 g, 10.7 mmol) in 50 mL of acetic acid was stirred at room temperature for 18 h. The solvent was

- 47 -

removed by rotoevaporation, and the residue was taken up in an diethyl ether/water mixture. The resulting solid was collected by filtration, and washed with diethyl ether. After drying the product (2.75 g) was
5 obtained in 95% yield.

Example 4

Preparation of acetic acid 8-acetylamino-5-bromo-7-nitro-1,2,3,4-tetrahydronaphthalen-2-yl ester




10

A solution of acetic acid 8-acetylamino-5-bromo-1,2,3,4-tetrahydronaphthalen-2-yl ester (2.6 g, 8 mmol) in 40 mL of trifluoroacetic acid was cooled in an ice bath. Fuming nitric acid (2 mL) was added dropwise and the mixture was stirred for 2.5 h. The solvent was removed by rotoevaporation, and water was added to the residue which induced solidification. The solid was collected by filtration, washed with ether and dried to
15 give the product (2.4 g) in 81% yield.
20

Example 5


Preparation of 8-amino-5-bromo-7-nitro-1,2,3,4-tetrahydronaphthalen-2-ol



5 A mixture of acetic acid 8-acetylamino-5-bromo-7-nitro-1,2,3,4-tetrahydronaphthalen-2-yl ester (2 g, 5.4 mmol), acetic acid (15 mL) and 6 N HCl (30mL) was
10 stirred at 90°C for 18 h. The solvent was removed by rotoevaporation to give the product (1.6 g) as an orange solid in quantitative yield.

Example 6


15 Preparation of 7,8-diamino-5-bromo-1,2,3,4-tetrahydronaphthalen-2-ol



20 Raney nickel was deactivated prior to use by washing twice with acetone, and then washing twice with tetrahydrofuran. A mixture of 8-amino-5-bromo-7-nitro-1,2,3,4-tetrahydronaphthalen-2-ol (1.8 g) and Raney
25 nickel (4 g) in tetrahydrofuran (100 mL) was stirred under a hydrogen atmosphere (1 atm) for 1 h. The catalyst was removed by filtration, and the filtrate evaporated. The solid residue was washed with diethyl ether and then dried to give the desired product.

Example 7


Preparation of 6-bromo-9-hydroxy-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione



A mixture 7,8-diamino-5-bromo-1,2,3,4-tetrahydronaphthalen-2-ol (0.5 g, 2 mmol) and oxalic acid (0.5 g, 4 mmol) in 2 N HCl (20 mL) was heated at 90°C for 3 h. A solid was collected by filtration, washed consecutively with water and diethyl ether and dried to give the quinoxalinedione product (0.33 g). CHN: calc'd for $C_{12}H_{11}BrN_2O_3 \cdot 1.5H_2O$ - C, 42.62; H, 4.17; N, 8.29; found - C, 42.87; H, 3.73; N, 8.29.

Example 8

Preparation of 9-benzyloxy-6-bromo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione



A mixture of 6-bromo-9-hydroxy-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione (0.1 g, 0.32 mmol) and sodium hydride (16 mg, 0.4 mmol) in dimethylformamide (1 mL) was stirred at 0°C for 30 min, and then treated with benzyl bromide (70 mg, 0.4 mmol). The reaction


- 50 -

mixture was stirred for 48 h and warmed to room temperature during that time. The solvent was removed by rotoevaporation, the residue triturated in ethyl acetate, and the solid collected by filtration and dried. CI MS m/e (M+1) 402, 404. H-NMR (200 MHz, dmsO):
1H, s, 12.05; 1H, s, 11.95; 5H, m, 7.4-7.2; 1H, s, 7.05; 2H, s, 5.4; 1H, s, 4.75; 1H, m, 3.9; + ring protons.

10

Example 9

Preparation of methanesulfonic acid
8-acetylamino-1,2,3,4-tetrahydronaphthalen-2-yl ester




15

A solution of N-(7-hydroxy-5,6,7,8-tetrahydronaphthalen-1-yl)-acetamide (5 g, 24 mmol), methylsulfonylchloride (6.9 g, 60 mmol) and pyridine (4.7 g, 60 mmol) in methylene chloride (100 mL) was heated at reflux for 8 h. The mixture was then cooled and washed with water. The organic layer was dried over magnesium sulfate, filtered and evaporated. The resulting solid was washed with diethyl ether and dried (3.8 g, 51% yield) without further purification.

25

Example 10


Preparation of N-(7-azido-5,6,7,8-tetrahydronaphthalen-1-yl)-acetamide



A mixture of methanesulfonic acid 8-acetylamino-1,2,3,4-tetrahydronaphthalen-2-yl ester (3.8 g, 13 mmol) and sodium azide (1.7 g, 27 mmol) in dimethylformamide (70 mL) was heated at 50°C for 6 h. The solvent was removed by rotoevaporation in vacuo and the residue partitioned between methylene chloride and water. The organic layer was separated, dried over sodium sulfate, filtered and evaporated to give a dark solid. The product was purified by column chromatography on silica gel (1:1 ethyl acetate:hexane as eluant) to give a white solid (2.9 g) in 97% yield.

Example 11


Preparation of N-(7-azido-4-bromo-5,6,7,8-tetrahydronaphthalen-1-yl)-acetamide



A solution of N-(7-azido-5,6,7,8-tetrahydronaphthalen-1-yl)-acetamide (1 g, 4.3 mmol) in acetic acid (40 mL) was treated dropwise with bromine (1 g, 6.5 mmol) and stirred at room temperature for 3 h. The solvent was evaporated and the solid residue was washed with ether, collected by filtration and dried to give the product (1.5 g) in 89% yield.

Example 12

Preparation of N-(7-azido-4-bromo-2-nitro-5,6,7,8-tetrahydronaphthalen-1-yl) acetamide




5

A solution of N-(7-azido-4-bromo-5,6,7,8-tetrahydronaphthalen-1-yl)-acetamide (1.5 g) was cooled in an ice bath and treated with fuming nitric acid (2 mL). After stirring for 2 h the solvent was removed by rotoevaporation and water was added to the residue to give a solid. The product was collected by filtration and dried (1.1 g, 81% yield).

10

Example 13

Preparation of 7-azido-4-bromo-2-nitro-5,6,7,8-tetrahydronaphthalen-1-ylamine



15


A mixture of N-(7-azido-4-bromo-2-nitro-5,6,7,8-tetrahydronaphthalen-1-yl)-acetamide, 2 N HCl (150 mL) and acetic acid (30 mL) was heated at 85°C for 48 h. The mixture was concentrated by rotoevaporation in vacuo, and the resulting solid was collected by filtration, washed with water and then dried to give the desired aniline (4.25 g) in 86% yield.

20

25

Example 14

Preparation of 9-amino-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione




5

A solution of 7-azido-4-bromo-2-nitro-5,6,7,8-tetrahydronaphthalen-1-ylamine (1.7 g, 5.4 mmol) in tetrahydrofuran (50 mL) and methanol (50 mL) was treated with 20% palladium on charcoal (0.5 g) and shaken in a Parr apparatus under a hydrogen atmosphere (50 psi) for 15 h. The catalyst was removed by filtration, the filtrate was evaporated and the residue was dissolved in 2 N HCl and treated with oxalic acid (1.7 g). After heating at 85°C for 2.5 h, the solid that formed was collected by filtration and washed with methanol and dried in vacuo to give the product (1.02 g, 82% yield).

20

Example 15

Preparation of 9-amino-6-nitro-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione



25


A mixture of 9-amino-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione and trifluoroacetic acid (10 mL) was cooled in an ice bath and then treated with

- 54 -

fuming nitric acid (0.5 mL) and stirred for 3 h at 0°C and then 1 h at room temperature. After removing the solvent by rotoevaporation, the residue was triturated with acetone and the resulting solid collected by filtration to give the product (0.22 g, 93% yield). The solid was stirred in 2 N HCl for 15 min, collected by filtration and dried to give the hydrochloride salt. Anal. calc'd for $C_{12}H_{12}N_4O_4 \cdot HCl \cdot H_2O$ - C, 43.58; H, 4.57; N, 16.94; found - C, 43.56; H, 4.08; N, 16.51.

Example 16

Preparation of piperidine-4-carboxylic acid
(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10
-octahydro-benzo[f]quinoxalin-9-yl) amide




N-t-butyloxycarbonylpiperidine-4-carboxylic acid (0.35 g, 1.5 mmol) and carbonyldiimidazole (0.24 g, 1.5 mmol) were refluxed in tetrahydrofuran (5 mL) for 15 min. The mixture was added to a solution of 9-amino-6-nitro-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione (0.15 g, 1.5 mmol) in dimethylformamide (5 mL) and heated to 70°C. After 15 min, triethylamine (0.15 g, 1.5 mmol) was added and heating continued at 70°C for 6 h. The solvent was removed by rotoevaporation in vacuo and the residue triturated with diethyl ether to form a solid. The solid was collected by filtration, washed with water and dried to give the crude product (0.23 g, 94% yield). For further purification, the solid was suspended and stirred in 2 N HCl, and then filtered and

- 55 -

dried. Anal. calc'd for $C_{23}H_{29}N_3O_7 \cdot 1.5 NaCl \cdot H_2O$: C, 40.82; H, 4.57; N, 13.22; found - C, 40.35; H, 4.70; N, 12.91.

Example 17

- 5 Preparation of N-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-octahydro-benzo[f]quinoxalin-9-yl)-acetamide




- 10 A mixture of 9-amino-6-nitro-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione (0.15 g, 0.5 mmol), acetic anhydride (0.1 g, 1 mmol) and triethylamine (0.2 g, 2 mmol) in dimethylformamide (5 mL) was stirred at room temperature for 3 h. The solvent was removed by
15 rotoevaporation, and the residue suspended in water and 2 N HCl. The solid which formed was collected by filtration, washed with diethyl ether and dried to give the acetamide (0.14 g) in 88% yield. Anal. calc'd for $C_{14}H_{14}N_4O_5 \cdot H_2O$: C, 50.00; H, 4.80; N, 16.66; found - C,
20 49.92; H, 4.77; N, 16.04.

Example 18

- Preparation of N-7-benzyl-4-bromo-2-nitro-5,6,7,8-tetrahydronaphthalene-1,7-diamine

25




- 56 -

A solution of 1,7-diamino-2-nitro-4-bromotetralin hydrochloride (0.32 g, 1 mmol) and benzaldehyde (0.16 g, 1.5 mmol) in 2:1 methanol:water (10 mL) was stirred at room temperature and treated with sodium cyanoborohydride (0.19 g, 3 mmol). After stirring for 18 h, the mixture was concentrated and extracted with methylene chloride. The organic layer was dried over magnesium sulfate, filtered and evaporated to give a yellow solid. Purification was accomplished by silica gel chromatography (chloroform, then 2% methanol in chloroform as eluant) to give the benzylamine derivative (0.18 g).

Example 19

Preparation of 9-benzylamino-6-bromo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione



Raney nickel was deactivated by stirring with acetone and washing with tetrahydrofuran as before. A mixture of N-7-benzyl-4-bromo-2-nitro-5,6,7,8-tetrahydronaphthalene-1,7-diamine (0.18 g) and Raney nickel (0.5 g) in tetrahydrofuran (20 mL) was stirred under a hydrogen atmosphere (1 atm) for 30 min. The catalyst was removed by filtration and washed with excess tetrahydrofuran. The filtrate was evaporated to a white solid. The catalyst was washed further with 2 N HCl and the filtrate was combined with the white solid from above and oxalic acid (0.6 g) and the mixture heated at 75°C for 2 h. The solid that formed was collected by filtration and dried in vacuo. CI MS


- 57 -

m/e (M+1) 400, 402. H-NMR (400 MHz, TFA): 1H, s, 12.05; 1H, s, 11.95; 1H, s, 7.7; 5H, s, 7.5; 1H, d, 4.8; 1H, d, 4.55; 1H, m, 3.95; 1H, m, 3.87; 2H, m, 3.2-2.4; 1H, m, 3.0-2.85; 1H, m, 2.6; 1H, m, 2.2-2.0.

5

Example 20

Preparation of N-indan-2-yl-acetamide




10

A mixture of 2-aminoindan hydrochloride (13.3 g, 78 mmol), acetic anhydride (6 g, 0.157 mmol) and saturated sodium bicarbonate (100 mL) in ether (100 mL) was stirred at room temperature for 1 h. The organic phase was separated, and the aqueous layer back washed with ether. The combined organic layer was washed with saturated chloride solution, dried over magnesium sulfate, filtered and evaporated to give the acetamide (13.6 g) in quantitative yield.

20

Example 21

Preparation of N-(5-nitro-indan-2-yl)-acetamide



25


N-Indan-2-yl-acetamide (14 g, 80 mmol) was dissolved in trifluoroacetic acid (150 mL) and cooled in an ice bath to 0°C. Fuming nitric acid (20 mL) was added slowly via a pipette, and the mixture was stirred for 2 h while maintaining the temperature at 0°. After evaporating the solvent in vacuo, the residue was dissolved in a mixture of ether and water. The water layer was extracted several times with ether, and the

30

combined organic layer was washed with saturated sodium chloride solution, dried over magnesium sulfate, filtered and evaporated to give the nitrated adduct(s) as a syrupy solid (20.5 g). This material (7:1 mixture
5 by NMR) was used without further purification.

Example 22

Preparation of N-(5-acetylamino-indan-2-yl)-acetamide




The mixture of N-(5-nitro-indan-2-yl)-acetamide (20.5 g, 93 mmol) in tetrahydrofuran (200mL) was combined with acetic anhydride (10mL) and Raney nickel (5 g) and
15 stirred under an atmosphere of hydrogen gas (in a balloon) at room temperature for 24 h. Additional Raney nickel was added and the balloon recharged with hydrogen gas and stirred an additional 24 h. After
20 removing the Raney nickel by filtration through a celite pad and washing with methanol several times, the filtrate was evaporated to a syrup. The residue was triturated with ether/water and the resulting white solid collected by filtration (9.2 g, 52% yield overall from 2-aminoindan, prepared in Example 20).

25

Example 23

Preparation of N-(5-acetylamino-6-bromo-indan-2-yl)-acetamide




A solution of N-(5-acetylamino-indan-2-yl)-acetamide (9.8 g, 42 mmol) and bromine (8.5 g, 53 mmol) in acetic acid (200 mL) was stirred at room temperature for 3 h.

- 59 -

After evaporation of the solvent, the syrupy residue was dissolved in ether and washed with water. The organic layer was washed with aqueous sodium bisulfite and dried over magnesium sulfate. The solvent was concentrated and the resulting solid collected by filtration (10 g, 77% yield).

Example 24


Preparation of N-(5-acetylamino-6-bromo-4-nitro-indan-2-yl)-acetamide



A solution of N-(5-acetylamino-6-bromo-4-indan-2-yl)-acetamide (10 g, 32 mmol) in trifluoroacetic acid (175 mL) was cooled in an ice bath to 0°C and then treated with fuming nitric acid (25 mL). After stirring for 2 h at 0°C, the reaction mixture was warmed to room temperature and stirred for an additional 2 h. The solvent was removed by rotoevaporation and the residue was triturated with an ether/water mixture. The resulting solid was collected by filtration and washed consecutively with water and ether, and then dried in vacuo to give the product (10 g) in 87% yield.

Example 25


Preparation of N-(5-amino-6-bromo-4-nitro-indan-2-yl)-acetamide



N-(5-Acetylamino-6-bromo-4-nitro-indan-2-yl)-acetamide
(1 g, 2.8 mmol) was suspended in a 2:1 mixture of
sulfuric acid and water (30 mL) and heated at 90°C for
12 h. The mixture was poured onto ice and the
5 resulting yellow solid was collected by filtration and
dried in vacuo to give (0.7 g, 80% yield).

Example 26

Preparation of 6-bromo-4-nitro-indan-2,5-diamine
10 monohydrochloride



N-(5-Acetylamino-6-bromo-4-nitro-indan-2-yl)-acetamide
15 (1 g, 2.8 mmol) was stirred in 3 N HCl (80 mL) at
reflux for 3 h. The resulting solid was collected by
filtration and dried in vacuo. Additional material was
obtained by evaporation of the filtrate to give a
combined yield of 93% of the yellow orange product (0.8
20 g).

Example 27

Preparation of N-(4,5-diamino-6-bromo-indan-2-yl)-
25 acetamide




N-(5-Amino-6-bromo-4-nitro-indan-2-yl)-acetamide (0.7
g, 2.2 mmol) was dissolved in tetrahydrofuran (20 mL)
30 and then treated with Raney nickel and stirred under a
hydrogen atmosphere (1 atm) at room temperature for 1
h. The catalyst was removed by filtration through a

- 61 -

celite pad, and washed repeatedly with methanol. The combined organic filtrate was removed by rotoevaporation to give a syrup that solidified upon standing. This product was used without further purification.

Example 28


Preparation of N-(5-bromo-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide



N-(4,5-Diamino-6-bromo-indan-2-yl)-acetamide (~2.2 mmol) was combined with oxalic acid (0.5 g) in 2 N HCl (15 mL) and heated at 80°C for 3 h. The resulting precipitate was collected by filtration and washed consecutively with methanol, water and ether, and then dried in vacuo to give the quinoxalinedione (0.37 g) in 50% yield for the hydrogenation and condensation steps. CHN calc'd for $C_{13}H_{12}BrN_3O_3 \cdot 2.2H_2O$: C, 41.33; H, 4.37; N, 11.12; Found: C, 41.12; H, 3.91; N, 11.24.

Example 29

Preparation of N-(5-bromo-6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide




- 62 -

N-(5-Bromo-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide (0.24 g, 0.7 mmol) was dissolved in trifluoroacetic acid (10 mL), cooled in an ice bath to 0°C, and then treated with fuming nitric acid (0.5 mL) and stirred for 4 h at 0°C. After removing the solvent by rotoevaporation, the syrupy residue was triturated with water and the resulting solid was collected by filtration, washed with water and then ether and dried in vacuo (0.22 g, 81% yield). Anal. calc'd for $C_{13}H_{11}BrN_4O_3 \cdot H_2O$: C, 38.92; H, 3.27; N, 13.94. Found: C, 38.60; H, 3.00; N, 13.66.

Example 30

Preparation of N-(2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide




A mixture of N-(5-amino-6-bromo-4-nitro-indan-2-yl)-acetamide (0.5 g, 1.6 mmol), 5% palladium on carbon (0.2 g) and sodium acetate (0.16 g) in methanol (75 mL) was placed on a Parr hydrogenation apparatus under a hydrogen atmosphere (51 psi) and shaken for 18 h. After removing the catalyst by filtration, the solvent was evaporated and the residue was suspended in 2 N HCl, treated with oxalic acid (0.5 g) and then heated at 80°C for 4 h. The water was evaporated and the residue dissolved in saturated sodium bicarbonate and acetic anhydride (1 mL) was added. The resulting solid was collected by filtration and dried in vacuo to give a brown solid (0.15 g) in 36% yield.

- 63 -

Example 31

Preparation of N-(2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide




5

N-(5-Bromo-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide (1 g, 3 mmol) was dissolved in dimethylformamide (75 mL) and treated with potassium acetate (0.33 g) and 20% palladium on carbon (0.2 g) and shaken on a Parr apparatus under a hydrogen atmosphere (51 psi) for 7 min. The catalyst was removed by filtration, the filtrate evaporated and the residue triturated in water. The resulting solid was collected by filtration and dried in vacuo to give the debrominated product (0.75 g) in 98% yield. Anal. Calc'd for C₁₃H₁₃N₃O₃•1.7 H₂O: C, 53.85; H, 5.70; N, 14.49. Found: C, 53.87; H, 5.04; N, 14.15.

20

Example 32

Preparation of N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide



25

A mixture of N-(2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide (0.15 g, 0.6 mmol), potassium nitrate (0.07 g, 0.7 mmol) and


- 64 -

sulfuric acid (5 mL) was stirred at room temperature for 3 h. After pouring onto ice, the aqueous solution was saturated with sodium chloride and allowed to stand overnight. The resulting precipitate was collected by filtration, washed with water and dried in vacuo to give a solid (35 mg, 19% yield). Anal. calc'd for $C_{13}H_{12}N_4O_3 \cdot 0.75 NaCl$: C, 44.85; H, 3.47; N, 16.09. Found: C, 45.18; H, 3.87; N, 15.87.

10

Example 33

Preparation of 8-amino-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione hydrochloride




15

N-2,3-Dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide (0.5 g, 1.9 mmol) was stirred in 2 N HCl (20 mL) at reflux for 72 h. The solid was collected by filtration and dried in vacuo (0.4 g, 83% yield).

20

Example 34

Preparation of 4-(2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-ylcarbamoyl)-piperidine-1-carboxylic acid tert-butyl ester




- 65 -

A mixture of N-Boc-piperidine-4-carboxylic acid (0.23 g, 1 mmol) and 1,1'-carbonyldiimidazole (0.16 g, 1 mmol) in tetrahydrofuran (5 mL) was heated at 80°C for 15 min. After cooling, it was added to a solution of
5 8-amino-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione hydrochloride (0.11 g, 0.5 mmol) and the mixture was heated at 70°C for 2 days. After filtration to remove precipitate, the filtrate was evaporated and the residue suspended in water. The
10 resulting solid was collected by filtration, washed with diethyl and dried in vacuo (0.15 g, 70% yield).

Example 35

Preparation of 4-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-ylcarbamoyl)-
15 piperidine hydrochloride



20 4-(2,3-Dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-ylcarbamoyl)-piperidine-1-carboxylic acid tert-butyl ester (0.15 g, 0.3 mmol) was dissolved in methylene chloride (5 mL) and cooled in an ice bath. Trifluoroacetic acid (0.2 mL) was added and
25 the mixture stirred for 20 min. The solvent was evaporated and the residue dissolved in trifluoroacetic acid (5 mL), cooled in an ice bath and treated with fuming nitric acid (0.5 mL), stirred for 2 h at 0°C and then warmed to room temperature for an additional 1 h.
30 The trifluoroacetic acid was removed by rotoevaporation, and the residue trituated with 2 N


- 66 -

HCl. The resulting solid was collected by filtration and dried in vacuo (60 mg, 42% yield).

Example 36

5

Preparation of N-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-octahydrobenzo[f]quinoxalin-9-yl)-benzamide.




- 10 Benzoic acid (0.15 g, 1.5 mmol) and carbonyldiimidazole (0.24 g, 1.5 mmol) were refluxed in 5 mL tetrahydrofuran for 15 min. The solution was added to the starting amine in dimethylformamide (5 mL) and heated to 70°C. Triethylamine was added after 30
- 15 min., and heated an additional 4h. The solvent was removed by rotoevaporation, and the residue was triturated with diethyl ether. The yellow solid was collected by filtration and dried in vacuo. CHN calc'd for C₁₈H₁₆N₂O₃·H₂O: C, 57.28; H, 4.55; N, 14.06; found - C, 57.21; H, 3.95; N, 14.00. CI MS m/e 398. H-NMR (400
- 20 MHz, dmsO): 1H, s, 12.1; 1H, s, 11.45; 1H, d, 8.55; 2H, d, 7.9, 1H, s, 7.7; 3 H, m 7.6-7.45; 1H, m, 4.3; + ring protons.

- 67 -

Example 37

Preparation of N-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-octahydrobenzo[f]quinoxalin-9-yl)cyclohexylamide.

5




Cyclohexanecarboxylic acid (0.13 g, 1 mmol) and carbonyldiimidazole (0.16 g, 1 mmol) were refluxed in 2 mL tetrahydrofuran for 15 min. The solution was added to the starting amine in dimethylformamide (2 mL) and heated to 60°C. Triethylamine was added after 30 min, and heated an additional 16h at 60°C. The solvent was removed by rotoevaporation, and the residue was taken up in 2N HCl. The yellow solid was collected by filtration and dried in vacuo to give 90 mg material. CHN calc'd for $C_{18}H_{22}N_4O_3 \cdot 0.75H_2O$: C, 57.06; H, 5.92; N, 14.01; found - C, 57.11; H, 5.50; N, 13.63. CI MS m/e 387. H-NMR (400 MHz, dmsO): 1H, s, 12.1; 1H, s, 11.45; 1H, d, 7.85; 1H, s, 7.7; 1H, m, 4.0; + ring protons.

Example 38

Preparation of 8-amino-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione hydrochloride.

25




The parent quinoxalinedione (Example 33) was mixed with 50 mL trifluoroacetic acid and 1 mL fuming nitric acid at 0°C and stirred for 3 h, then warmed to room temperature for 2 h. The solvent was evaporated, the residue taken up in acetone, and the solid collected by filtration. After washing with water and then tetrahydrofuran, the material was dried to give 1.3 g of the compound as the hydrochloride salt. CHN calc'd for $C_{11}H_{10}N_4O_4 \cdot HCl \cdot 0.5H_2O$; C, 38.39; H, 3.52; N, 16.28; found - C, 38.54; H, 3.57; N, 17.39. CI MS m/e (M+1) 263. H-NMR (400 MHz, dmsO) 1H, s, 12.2; 1H, s, 12.12; 2H, br s, 8.1; 1H, s, 7.9; 1H, br s, 4.2; 1H, dd, 3.7; 2H, m, 3.4; 1H dd, 3.1.

15

Example 39

Preparation of N-(6-nitro-2,3-dioxo-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxalin-8-yl)-benzamide.



20


The product of Example 38 (0.13 g, 0.5 mmol), benzoic anhydride (0.23g, 1 mmol) and triethylamine (0.1 g) were combined in 5 mL dimethylformamide and stirred at room temperature for 24 h. The solvent was evaporated, the residue washed with methanol, collected by filtration and dried to give 0.1 g. CHN calc'd for $C_{18}H_{14}N_4O_5 \cdot H_2O$; C, 56.24; H, 4.20; N, 14.58; found - C, 56.33; H, 3.82; N, 14.32. CI MS m/e 367. H-NMR (400 MHz, dmsO) 1H, s, 12.18; 1H, s, 11.95; 1H, d, 8.7; 3H, m, 7.82; 3H, m, 7.55-7.4; 1H, m, 4.8; + ring protons.

30

- 69 -

Example 40


Preparation of 4-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-octahydrobenzo[f]quinoxaline-9-ylcarbamoyl)-piperidine hydrochloride.



A mixture of N-Boc-piperidine-4-carboxylic acid (0.92 g, 4 mmol) and carbonyldiimidazole (0.65 g, 4 mmol) in 10 mL tetrahydrofuran was heated at reflux for 15 min. The mixture was added to the starting amine (Example 15, 0.6 g, 2 mmol) in 10 mL dimethylformamide and heated at 70°C for 30 min. Triethylamine (0.4 g) was added and the mixture heated an additional 18 h. The solvent was evaporated and the yellow solid was washed with tetrahydrofuran, collected by filtration and dried to give 0.74g. The solid was suspended in 50 mL methylene chloride, cooled in an icebath and treated with trifluoroacetic acid (1 g). After evaporating the solvent, the solid was washed with ether and acetone mixture and dried. Calc'd for $C_{18}H_{21}N_5O_5 \cdot 1.5HCl \cdot H_2O$: C, 40.82; H, 4.57; N, 13.22; found - C, 40.35; H, 4.70; N, 12.91.

Example 41

Preparation of 2-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-octahydrobenzo[f]quinoxaline-9-ylcarbamoyl)ethyl-(4-hydroxy)benzene.




Hydroxyphenylpropionic acid (0.17 g, 1 mmol) and carbonyldiimidazole (0.16 g, 1 mmol) in 5 mL tetrahydrofuran were stirred at reflux for 15 min. The mixture was added to the starting amine (example 15, 0.15 g, 0.5 mmol) and heated at 60°C for 30 min. Triethylamine (0.1 g) was added and the mixture heated an additional 5 h. The solvent was removed by rotoevaporation, and the residue suspended in tetrahydrofuran. The solid was collected by filtration and dried. CI MS m/e (M+1) 425. H-NMR (400 MHz, dmsO) 1H, s, 12.1; 1H, s, 11.45; 5 Ar H, 8.0-6.6 including 1 H, s, 7.65; 1 H, m, 4.1; + other ring protons.

- 71 -

Example 42

Preparation of N-phenyl-N'-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-octahydrobenzo[f]quinoxalin-9-yl)-urea.

5



The product of Example 15 (0.15 g, 0.5 mmol), phenylisocyanate (0.1 g, 1 mmol) and triethylamine (0.1g) in 5 mL dimethylformamide were stirred at room temperature for 18 h. The solvent was removed by rotoevaporation and the residue suspended in methanol/dimethylformamide. After heating, the solid was collected by filtration and dried in vacuo. CHN calc'd for $C_{19}H_{17}N_3O_3 \cdot 0.5H_2O \cdot HCON(Me)_2$: C, 55.34; H, 5.27; N, 17.60 found - C, 55.38; H, 4.95; N, 17.04. H-NMR (400 MHz, dmsO) 1 H, s, 12.1; 1 H, s, 11.45; 1H, s, 8.4; 1H, s, 7.95; 1H, s, 7.7; 2H, d, 7.4; 2H, t, 7.23; 1H, t, 6.9 1H, d, 6.35; 1H, t, 4.1; + ring protons.

20

Example 43


8-Acetamido-2-methoxynaphthalene



5 A mixture of 8-Acetamido-2-naphthol (21 g, 0.104 mol),
dimethyl sulfate (15.1 g, 0.12 mol) and potassium
carbonate (41 g, 0.3 mol) in acetone (200 mL) was
stirred at room temperature for 48 h. Solids were
10 removed by filtration and washed with methanol, and the
filtrate evaporated. The residue was washed in 1:1
hexane:toluene, collected by filtration and dried (24.7
g).

Example 44

8-Acetamido-2-tetralone



20 Ammonia (300 mL) was condensed in a flask containing 8-
Acetamido-2-methoxynaphthalene (24.5 g, 115 mmol), t-
butyl alcohol (34 g, 0.46 mol) and THF (300 mL).
Sodium (8 g, 0.35 mol) was added in portions and the
mixture was stirred for 3 h and allowed to warm to room
25 temperature. The mixture was poured onto ice and
saturated sodium chloride, and then extracted with THF
(3x). The combined organic layers were washed with
saturated sodium chloride, dried over magnesium
sulfate, filtered and evaporated to give a dark solid.
30 The residue was dissolved in 50% acetic acid and heated
at 95°C for 18 h. After evaporation, the residue was
suspended in dimethyl ether and collected by filtration


- 73 -

and dried. The filtrate was purified by silica gel chromatography (3:2 EtOAc:hexane) and combined with the other product to give a tan solid (10 g).

5

Example 45


1-Acetamido-7-benzylamino-5,6,7,8-tetrahydronaphthalene



- 10 A solution of 8-acetamido-2-tetralone (15.5 g, 76 mmol), benzylamine (16.3 g, 150 mmol) and tosic acid (0.2 g) in benzene (200 mL) and DMF (50 mL) was refluxed with a Dean-Stark trap attached for 18 h. The solvent was evaporated and the residue dissolved in 1:1
- 15 methanol:THF (250 mL). After cooling in an ice bath, sodium cyanoborohydride was added and the mixture warmed to room temperature and stirred for 5 h. The mixture was basified with 12.5% NaOH and the solvent concentrated. The residue was dissolved in methylene
- 20 chloride:water and filtered through a celite pad. The aqueous layer was extracted with methylene chloride, and the combined organic layers were dried over sodium sulfate, filtered and evaporated. The dark residue was purified by column chromatography (9:1 EtOAc:methanol)
- 25 to give the product (15 g).

Example 46


1-Acetamido-7-benzylmethylamino-5,6,7,8-tetrahydronaphthalene



A mixture of 1-acetamido-7-benzylamino-5,6,7,8-tetrahydronaphthalene (14.3 g, 50 mmol), paraformaldehyde (15 g, 0.5 mol) and sodium cyanoborohydride (15.7 g, 0.25 mol) in acetic acid (250 mL) was stirred at room temperature for 48 h. The solvent was removed and the residue taken up in water:diethyl ether. The aqueous phase was basified with 12.5% NaOH and back extracted with ether. The combined ether layers were washed with saturated NaCl, dried over magnesium sulfate, filtered and evaporated to give a viscous syrup (16 g).

Example 47

20 1,7-Bis(N⁺-methyl)acetamido-5,6,7,8-tetrahydronaphthalene



25 A mixture of 1-acetamido-7-benzylmethylamino-5,6,7,8-tetrahydronaphthalene (14 g, 45 mmol) and 20% Pd/C (3 g) in acetic acid (200 mL) was shaken on a Parr hydrogenation apparatus under a hydrogen atmosphere (50 psi) for several hours (hydrogen replaced periodically). After removing the catalyst the solvent

30


- 75 -

was evaporated and the residue portioned between methylene chloride and water. The aqueous layer was basified with 12.5% NaOH and back extracted. The combined organic layers were dried over sodium sulfate, filtered and evaporated. The residue was purified by column chromatography (gradient from 30 - 100% methanol in EtOAc) to give the methylamine adduct (7.4 g) as a clear solid. The methylamine (5 g, 23 mmol) was dissolved in THF (50 mL) and treated with acetic anhydride (6 g, 60 mmol) and the mixture stirred at room temperature for 1 h. The solvent was removed, and the residue triturated with dimethyl ether and the solid collected by filtration and dried (5.6 g).

15

Example 48

1,7-Bis(N⁷-methyl)acetamido-4-bromo-5,6,7,8-tetrahydronaphthalene




20

A solution of 1,7-bis(N⁷-methyl)acetamido-5,6,7,8-tetrahydronaphthalene (7 g, 27 mmol) and sodium acetate trihydrate (5.5 g, 51 mmol) in acetic acid (100 mL) was treated with bromine (5.2 g, 32 mmol) and stirred at room temperature for 18 h. Sodium bisulfite was added and the solvent removed. The residue was taken up in water:methylene chloride and the aqueous phase back extracted. The combined organic layers were evaporated to give an off-white solid. This was washed with dimethyl ether and dried to give the product (8.5 g) as a white solid.

30

Example 49


1,7-Bis(N⁷-methyl)acetamido-4-bromo-2-nitro-5,6,7,8-tetrahydronaphthalene



A mixture of 1,7-bis(N⁷-methyl)acetamido-4-bromo-5,6,7,8-tetrahydronaphthalene (8.5 g, 25 mmol) in trifluoroacetic acid (100 mL) was cooled in an ice bath and treated with fuming nitric acid (10 mL). The reaction mixture was warmed to room temperature after 1 h and stirred for 3 h longer. The solvent was removed and the residue taken up in methylene chloride:water. The aqueous layer was backed extracted and the combined organic layers were dried over sodium sulfate, filtered and evaporated to give the product as a yellow syrup (9.2 g).

Example 50

20 1,7-Di(N⁷-methyl)amino-4-bromo-2-nitro-5,6,7,8-tetrahydronaphthalene hydrochloride




25 1,7-Di(N⁷-methyl)amino-4-bromo-2-nitro-5,6,7,8-tetrahydronaphthalene was hydrolyzed in 3 N HCl (100 mL) and acetic acid (20 mL) at 100°C for 72 h. The mixture was cooled in an ice bath for 1 h, and the orange precipitate collected by filtration and dried (7.6 g).

30

- 77 -

Example 51


9-Methylamino-1,4,7,8,9,10-hexahydro-
benzo[f]quinoxaline-2,3-dione hydrochloride



A mixture of 1,7-di(N^m-methyl)amino-4-bromo-2-nitro-5,6,7,8-tetrahydronaphthalene hydrochloride (7.6 g, 23 mmol) and 20% Pd/C (1 g) in methanol (250 mL) was shaken on a Parr hydrogenator under a hydrogen atmosphere (52 psi) for 18 h. After removing the catalyst, the solvent was removed and the residue dissolved in 2 N HCl (100 mL). Oxalic acid (6.3 g, 50 mmol) was added and the reaction heated at 100°C for 3 h. After cooling to room temperature, the precipitate was collected by filtration and dried.

Example 52

9-Methylamino-6-nitro-1,4,7,8,9,10-hexahydro-
benzo[f]quinoxaline-2,3-dione



9-Methylamino-1,2,3,4,7,8,9,10-octahydro-
benzo[f]quinoxaline-2,3-dione hydrochloride (1 g, 3.5 mmol) was nitrated using fuming nitric acid (1 mL) and trifluoroacetic acid (30 mL) from 0°C to room


- 78 -

temperature for 1 h. The solvent was removed and their residue triturated in acetone:water and collected by filtration and dried to give the product (1.1 g).
Calc'd for $C_{13}H_{14}N_4O_4 \cdot HCl$: C, 47.79; H, 4.63; N, 17.15;
5 found: C, 43.36; H, 4.22; N, 18.85. MS $M+1$ (291).

Example 53

N-Methyl-N-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-octahydro-benzo[f]quinoxalin-9-yl)-benzamide

10




A mixture of 9-Methylamino-6-nitro-1,2,3,4,7,8,9,10-octahydrobenzo[f]quinoxaline-2,3-dione (0.16 g, 0.5
15 mmol), benzoic anhydride (0.22 g, 1 mmol) and triethylamine (0.1 g, 1 mmol) in DMF (5 mL) was stirred at room temperature for 18 h. The solvent was removed, and the residue suspended in 2 N HCl. The solid was collected by filtration, washed successively with water
20 and dimethyl ether and dried. Calc'd for $C_{20}H_{18}N_4O_5 \cdot H_2O$: C, 59.81; H, 4.72; N, 13.95; found: C, 59.98; H, 4.60; N, 13.60.

- 79 -

Example 54

Thiophene-2-carboxylic acid (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-yl)-amide.

5




A solution of thiophene-2-carboxylic acid (1 mmol) in anhydrous tetrahydrofuran (THF) was treated with carbonyldiimidazole (1.2 mmol) and triethyl amine (1 mL) and heated at 60°C for 30 min. After cooling to room temperature a solution of 8-amino-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione (1 mmol) in dimethylformamide (DMF) was added. The reaction mixture was stirred at room temperature for 2 h, then heated at 70°C for 2 h. The solvent was evaporated and the residue triturated with acetone. The solid was collected by filtration, washed with diethyl ether and dried in vacuo. Calc'd for $C_{16}H_{12}N_4O_5S \cdot H_2O$: C, 49.22; H, 3.61; N, 14.35, found: C, 49.06; H, 3.12; N, 15.05. MS $M+1$ (373).

25

Example 55

Resolution of 2,5-diamino-6-bromo-4-nitroindane




- 80 -

The amine (10.1 g) was dissolved in 185 mL of 4:1 isopropanol:water. (R)-Mandelic acid (2.8 g, 0.5 eq) was added and the mixture allowed to crystallize at room temperature for 18 h. The crystals were collected by filtration and dried. The solid was partitioned between methylene chloride: 1 N NaOH, and the aqueous layer was back extracted. The combined organic layers were dried over sodium sulfate, and evaporated to give a solid (3.2 g). The recrystallization with (R)-mandelic acid was repeated a total of four times to give an enantiomeric excess of 99% of (R)-2,5-diamino-6-bromo-4-nitroindane as determined by chiral HPLC. Absolute configuration was determined by x-ray crystallography. The (R)- and (S)-enantiomers were carried individually through the remaining synthesis to give chiral quinoxalinedione adducts.

Example 56

(S)-2,5-diamino-6-bromo-4-nitroindane

20




The title compound was obtained by an identical procedure to Example 55 except that the racemate was cocrystallized with (S)-mandelic acid.

25

Example 57

(R) - Thiophene-2-carboxylic acid
 (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-
 1H-cyclopenta[f]quinoxalin-8-yl) -amide

5




(S) -2,5-diamino-6-bromo-4-nitroindane was hydrogenated
 to the desbromotriamine and then condensed with oxalic
 10 acid (3 N HCl at reflux) to give the 8-amino-4,7,8,9-
 tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione. This
 intermediate was nitrated at C-6 in a solution of
 trifluoroacetic acid and fuming nitric acid at a
 temperature of 0°C and then warmed to room temperature.
 15 The thienyl amide title compound was formed in an
 identical manner as described in example 54.

Example 58

(S)-Thiophene-2-carboxylic acid
(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-
1H-cyclopenta[f]quinoxalin-8-yl)-amide

5




The title compound was prepared from (R)-2,5-diamino-6-bromo-4-nitroindane in a manner similar to that described in Example 57.

10

Example 59

Furan-3-carboxylic acid (6-nitro-2,3-dioxo-
2,3,4,7,8,9-hexahydro-1H-
cyclopenta[f]quinoxalin-8-yl)-amide

15



A solution of furan-3-carboxylic acid (1 mmol) in anhydrous THF was treated with carbonyldiimidazole (1.2 mmol) and triethyl amine (1 mL) and heated at 60°C for 30 min. After cooling to room temperature a solution of 8-amino-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione (1 mmol) in DMF was added. The reaction mixture was stirred at room temperature for 2 h, then heated at 70°C for 2 h. The solvent was evaporated and the residue triturated with acetone. The solid was collected by filtration, washed with diethyl ether and dried in vacuo. Calc'd for

20


25

- 83 -

$C_{16}H_{12}N_4O_6 \cdot H_2O$: C, 51.34; H, 3.77; N, 14.97; found: C, 50.97; H, 3.18; N, 14.89. MS $M+1$ (357).

Example 60

5 Thiophene-3-carboxylic acid (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)amide



10


The title compound was prepared by a method identical to that used in example 54 except thiophene 3-carboxylic acid was used as a reactant. Calc'd for $C_{16}H_{12}N_4O_5S$: C, 51.61; H, 3.25; N, 15.05; found: C, 48.04; H, 3.45; N, 15.87. MS $M+1$ (373).

15

Example 61

N-(6-Nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-nicotinamide

20



The title compound was prepared by a method identical to that used in example 54 except that nicotinic acid was used as a reactant. Calc'd for $C_{17}H_{13}N_5O_5$: C, 55.59; H, 3.57; N, 19.07; found: C, 50.27; H, 3.56; N, 18.70. MS $M+1$ (368).


25

- 84 -

Example 62

Pyridine-2-carboxylic acid (6-nitro-2,3-dioxo-
2,3,4,7,8,9-hexahydro-
1H-cyclopenta[f]quinoxalin-8-yl)-amide


5



The title compound was prepared by a method identical
to that used in example 54 except that pyridine-2-
10 carboxylic acid was used as a reactant. Calc'd for
 $C_{17}H_{13}N_3O_5 \cdot H_2O$: C, 52.98; H, 3.92; N, 18.17; found: C,
52.70; H, 3.76; N, 18.29.

Example 63

15 2,3,4,5-Tetrahydroxy-tetrahydro-furan-2-carboxylic acid
(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-
1H-cyclopenta[f]quinoxalin-8-yl)-amide




20

The title compound was prepared by a method identical
to that used in example 54 except that furan-2-
carboxylic acid was used as a reactant. Calc'd for
 $C_{16}H_{16}N_4O_{10} \cdot H_2O$: C, 40.19; H, 4.35; N 13.39; found: C,
25 40.11; H, 4.50; N, 13.41.

Example 64

Pyrrolidine-2-carboxylic acid (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-amide

5




The title compound was prepared by a method identical to that used in example 54 except that pyrrolidine-2-carboxylic acid was used as a reactant. Calc'd for $C_{16}H_{17}N_3O_5 \cdot C_2H_5F_3O_2$: C, 45.67; H, 3.83; N, 14.79; found: C, 40.70; H, 3.66; N, 12.43.

15

Example 65

Benzo[b]thiophene-2-carboxylic acid (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-amide



20


The title compound was prepared by a method identical to that used in example 54 except that benzo[b]thiophene-2-carboxylic acid was used as a reactant. Calc'd for $C_{20}H_{14}N_4O_3S$: C, 56.87; H, 3.34; N, 13.26; found: C, 50.99; H, 3.78; N, 14.38. MS M+1 (423).

25

Example 66

2-(1H-Indol-3-yl)-N-(6-nitro-2,3-dioxo-
2,3,4,7,8,9-hexahydro-
1H-cyclopenta[f]quinoxalin-8-yl)-acetamide

5




The title compound was prepared by a method identical
to that used in example 54 except that 3-indolyl acetic
acid was used as a reactant. Calc'd for $C_{17}H_{17}N_3O_3 \cdot H_2O$:
C, 57.66; H, 4.38; N, 16.01; found: C, 57.56; H, 4.28;
N, 15.67.

15

Example 67

Thiophene-2-carboxylic acid methyl-
(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-
1H-cyclopenta[f]quinoxalin-8-yl)-amide



20


The title compound was prepared by a method identical
to that used in example 54 except that 8-(N-
methyl)amino-6-nitro-4,7,8,9-tetrahydro-1H-
cyclopenta[f]quinoxaline-2,3-dione was used as a
reactant. Calc'd for $C_{17}H_{14}N_2O_3S \cdot 0.75 H_2O$: C, 51.05; H,
3.91; N, 14.01; found: C, 50.81; H, 3.60; N, 13.48. MS
M+1 (387).

25

- 87 -

Example 68

N-Methyl-N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-benzamide



5


The title compound was prepared by a method identical to that used in example 67 except that benzoic acid was used as a reactant. Calc'd for $C_{19}H_{16}N_4O_5 \cdot 0.5 H_2O$: c, 58.60; H, 4.40; N, 14.39; found: C, 58.62; H, 4.35; N, 13.26. MS M+1 (381).

10

Example 69

6-Nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-carboxylic acid methyl-phenyl-amide

15



A solution of 6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-carboxylic acid (1 mmol, example 78) in anhydrous DMF was treated with carbonyldiimidazole (2 mmol) and triethylamine (1 mL) and then heated at 70°C for 45 min. After cooling to room temperature methylphenylamine (1.2 mL) was added and the mixture stirred at 50°C for 6 h. The solvent was removed in vacuo, and the residue triturated with


25

acetone. The solid was collected by filtration, washed with diethyl ether and dried in vacuo. Calc'd for $C_{19}H_{16}N_4O_3 \cdot 1.2 H_2O$: C, 56.77; H, 4.61; N, 13.93; found: C, 56.22; H, 4.13; N, 14.62. MS M+1 (381).

5

Example 70

2-Carboxyethyl-1-indanone



10

A mixture of 1-indanone (26.4 g, 0.2 mol), diethyl carbonate (35.4 g, 0.3 mol) and sodium hydride (16 g, 0.4 mol, 60% in oil) in THF (600 mL) was stirred at 40° C for 2 h. The mixture was then poured onto 500 mL of 2 N HCl and ice. The organic phase was separated, and the aqueous phase extracted with ether. The combined organic layers were washed with saturated NaCl, dried over Magnesium sulfate, filtered and evaporated to give the product as a syrup (38.3 gm).

20

Example 71

2-Carboxyethylindane



25

A mixture of 2-carboxyethyl-1-indanone (38.3 gm, 0.19 mol), 20% Pd/C (4 gm) and concentrated sulfuric acid (4 drops) in ethanol (400 mL) was shaken on a Parr hydrogenation apparatus under a hydrogen atmosphere (50.5 psi) for 15 h. After removal of the catalyst by filtration, the solvent was evaporated and the residue

30

purified by silica gel chromatography (4:1 hexane:EtOAc) to give the product (28.8 gm).

Example 72

5

2-Carboxyethyl-5-nitroindane



10 A mixture of 2-carboxyethylindane (28.8 g, 0.15 mol) and trifluoroacetic acid (300 mL) was cooled in an ice bath and then treated with fuming nitric acid (50 mL). The reaction mixture was slowly warmed to room temperature and allowed to stir for 18 h. After removing the solvent, the residue was dissolved in an
15 ether:water mixture. The ether layer was separated and washed with water, and then saturated NaCl, and dried over magnesium sulfate. The filtrate was evaporated to give a dark syrup (34.3 gm, 3:1 ratio of isomers).

20

Example 73

5-Acetamido-2-carboxyethylindane



25

A mixture of 2-carboxyethyl-5-nitroindane (34 g, 0.15 mol), Raney nickel (5 g), acetic anhydride (25 mL) and acetic acid (225 mL) was shaken on a Parr hydrogenation apparatus under a hydrogen atmosphere (50 psi) for 30
30 h. After removing the catalyst, the filtrate was evaporated to give a dark syrup. The residue was purified by column chromatography (3:2 hexane:EtOAc).

- 90 -

Example 74


5-Acetamido-6-bromo-2-carboxyethylindane



5 A solution of 5-acetamido-2-carboxyethylindane (18 g, 73 mmol), sodium acetate (9 g, 110 mmol) in acetic acid (300 mL) was treated dropwise with bromine (14.1 g, 88 mmol) and then stirred for 18 h. After removing the
10 solvent, the residue was dissolved in an ether:water mixture, and sodium bisulfite was added. The ether layer was separated and the aqueous layer back extracted. The combined organic layers were washed
15 with saturated NaCl, dried over magnesium sulfate, filtered and evaporated. The syrup was purified by silica gel chromatography (3:2 hexane:EtOAc) to give the product.

Example 75


20 5-Acetamido-6-bromo-2-carboxyethyl-4-nitroindane



25 A mixture of 5-acetamido-6-bromo-2-carboxyethylindane (7 g, 20 mmol) and trifluoroacetic acid (150 mL) was cooled in an ice bath and treated with fuming nitric acid (10 mL). After 2 h at 0° C, the solvent was removed and the residue solidified upon addition of water. The solid was collected by filtration, washed
30 with ether and dried. Crystallization from hot toluene gave the pure product (7 g).

Example 76


5-Amino-6-bromo-4-nitroindane-2-carboxylic acid



5 A mixture of 5-acetamido-6-bromo-2-carboxyethyl-4-nitroindane (7.1 g, 19 mmol) in acetic acid (25 mL) and 3 N HCl (100 mL) was heated at 100° C for 18 h. The mixture was concentrated by rotoevaporation, and the
10 resulting orange solid was collected by filtration, washed with ether and dried to give the product (5.5 g).

Example 77

2,3-Dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-carboxylic acid




20 A mixture of 5-amino-6-bromo-4-nitroindane-2-carboxylic acid (5.5 g, 18 mmol) and 20% Pd/C (0.5 g) in 95% ethanol (250 mL) and tetrahydrofuran (250 mL) was shaken on a Parr hydrogenation apparatus under a hydrogen atmosphere (51 psi) for 30 h. After removing
25 the catalyst, the solvent was evaporated to give a solid. The solid was dissolved in 2 N HCl (150 mL) and treated with oxalic acid (4.6 g), and then the mixture was heated at 90° C for 5 h. After cooling to room temperature, the precipitate was collected by
30 filtration and dried in vacuo (4.0 g).

- 92 -

Example 78

6-Nitro-2,3,-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-carboxylic acid



5

A mixture of 2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-carboxylic acid (4 g, 16 mmol) in trifluoroacetic acid (75 mL) was cooled to 0° C and then treated with fuming nitric acid (5 mL). After stirring in an ice bath for 3 h, the mixture was warmed to room temperature for 1 h, the solvent evaporated, and the residue was suspended in water. A solid precipitate was collected by filtration, washed with ether and dried in vacuo (4 g). Calc'd for $C_{15}H_8N_2O_6 \cdot 0.5 H_2O$: C, 48.00; H, 3.36; N, 14.00; found: C, 48.64; H, 3.13; N, 13.86. MS M+1 (292).

10

15

Example 79

20

General Procedure for Amide Preparation

A mixture of 6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-carboxylic acid (0.29 g, 1 mmol), the coupling reagent N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (EDAC) (0.29 g, 1.5 mmol), and the 'generic amine' (1.5 mmol) in dimethylformamide (15 mL) was treated with dimethylaminopyridine (10 mg) and heated and 50° C for 18 h. The solvent was evaporated, the residue taken up in water treated as necessary to cause precipitation of the product. The product was purified using standard methodologies. For example, aniline was couple by a similar method with 6-nitro-2,3-dioxo-2,3,4,7,8,9-


25

30

hexahydro-1H-cyclopenta[f]quinoxaline-8-carboxylic acid
to give the carboxylic acid phenylamide, example 57,
and N-methylaniline was coupled to give example 55.
Other couple reagents including carbonyldiimidazol
5 (CDI), dicyclohexylcarbodiimide (DCC), etc. can be
substituted for EDAC.

Example 80

10 6-Nitro-2,3,-dioxo-2,3,4,7,8,9-hexahydro-1H-
cyclopenta[f]quinoxaline-8-carboxylic acid phenylamide




The title compound was prepared by a method identical
15 to that used in example 69 except that aniline was used
as a reactant. Calc'd for $C_{18}H_{14}N_4O_3 \cdot H_2O$: C, 56.24; H,
4.20; N, 14.58; found: C, 56.10; H, 4.04; N, 14.57. MS
M+1 (367)

20

Example 81

1-Acetamido-7-pyrrolidin-1-yl-5,6,7,
8-tetrahydronaphthalene



25

A mixture of 8-acetamido-2-tetralone (0.4 g, 2 mmol),
pyrrolidine (0.28 g, 4 mmol) and tosic acid (10 mg) in
benzene:DMF (5:1, 12 mL) was heated at reflux with a

- 94 -


Dean-Stark trap for 4 h. The solvent was evaporated and the residue dissolved in 1:1 MeOH:THF (10 mL). Acetic acid (1 mL) and sodium cyanoborohydride (0.25 g, 4 mmol) were added, and the reaction stirred for 16 h.

5 The solvent was evaporated and the residue taken up in water:methylene chloride. The aqueous layer was basified with 12.5% NaOH, and back extracted. The combined organic phase was dried over sodium sulfate, filtered and evaporated. The residue solidified upon

10 standing, and was crystallized from toluene:hexane to give the product (0.36 g).

Example 82

15 1-Acetamido-4-bromo-7-pyrrolidin-1-yl-5,6,7,8-tetrahydronaphthalene



A solution of 1-acetamido-7-pyrrolidin-1-yl-5,6,7,8-tetrahydronaphthalene (0.36g, 1.4mmol) and sodium acetate trihydrate (0.38g, 2.8mmol) in acetic acid (15mL) was treated with bromine (0.27g, 1.7mmol) and stirred at room temperature for 2 h. A precipitate was collected by filtration and the filtrate evaporated.

20 The two solids were combined and dissolved in methylene chloride; water. The mixture was basified with 12.5% NaOH, and the organic phase separated. The aqueous layer was back extracted and the combined organic layers were dried over sodium sulfate, filtered and


25 evaporated. The residue was heated in diethyl ether to form a solid, which was collected and dried (0.31g).

30

- 95 -

Example 83

1-Acetamido-4-bromo-2-nitro-7-pyrrolidin-1-yl-
5,6,7,8-tetrahydronaphthalene




A mixture of 1-acetamido-4-bromo-7-pyrrolidin-1-yl-
5,6,7,8-tetrahydronaphthalene (0.3 g) in
trifluoroacetic acid (10 mL) was cooled in an icebath
10 and treated with fuming nitric acid (0.5 mL). After
stirring for 4 h, the solvent was evaporated and the
residue taken up in methylene chloride:water. The
mixture was basified with 12.5% NaOH, and the aqueous
layer back extracted. The combined organic layers were
15 dried over sodium sulfate, filtered and evaporated to
give a solid. Preparative thin layer chromatography
(7:3 EtOAc:MeOH) was used to obtain the purified
product (0.12 g).

20

Example 84

1-Amino-4-bromo-2-nitro-7-pyrrolidin-1-yl-5,6,7,
8-tetrahydronaphthalene hydrochloride




The acetamide (0.12 g) was hydrolyzed at 100°C in 2 N
HCl while stirring for 18 h. The solvent was removed
to give a yellow solid (0.1 g).

30

Example 85

9-Pyrrolidin-1-yl-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3,-dione hydrochloride

5



10 A mixture of 1-amino-4-bromo-2-nitro-7-pyrrolidin-1-yl-5,6,7,8-tetrahydronaphthalene hydrochloride (0.1 g, 0.27 mmol) and 20% Pd/C (0.03 g) in methanol (75 mL) was shaken on a Parr hydrogenation apparatus under a hydrogen atmosphere (51 psi) for 3 h. The solvent was evaporated and the residue dissolved in 2 N HCl (5 mL).


15 Oxalic acid (0.12 g) was added and the mixture heated at 100°C for 3 h. After cooling to room temperature, the precipitate was collected by filtration and dried (0.04 g).

- 97 -

Example 86

6-Nitro-9-pyrrolidin-1-yl-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione


5



A mixture of 9-pyrrolidin-1-yl-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione hydrochloride (35 mg) in trifluoroacetic acid (2 mL) was cooled in an ice bath and treated with 0.5 mL fuming nitric acid. After 2 h at room temperature, the solvent was evaporated and the residue suspended in acetone:water, and the resulting solid collected by filtration (35 mg). Calc'd for $C_{16}H_{18}N_4O_4 \cdot HCl$: C, 52.39; H, 5.22; N, 15.27; found: C, 42.95; H, 4.69; N, 16.28. MS $M+1$ (331).

Example 87

20 2,5-Di-(N²-dimethyl)amino-6-bromo-4-nitroindane



A mixture of 2,5-diamino-6-bromo-4-nitroindane hydrochloride (1.5 g, 5 mmol), paraformaldehyde (0.9 g, 10 mmol), sodium cyanoborohydride (0.63 g, 10 mmol) and sodium acetate (1.3 g, 10 mmol) in acetic acid was stirred at room temperature for 72 h, then heated at 50°C for 6 h. The solvent was evaporated and the residue dissolved in water:methylene chloride. The

- 98 -

aqueous layer was basified with 12.5% NaOH and back extracted. The combined organic layers were dried over sodium sulfate, filtered and evaporated to give a yellow solid. Silica gel chromatography (30% methanol in EtOAc) gave the product as an orange solid (0.5 g).

Example 88

2,5-Di-(N²-diethyl)amino-6-bromo-4-nitroindane


10

The title compound was prepared from 2,5-diamino-6-bromo-4-nitroindane hydrochloride (1.4 g, 5 mmol) and acetaldehyde (4.3 g, 0.1 mol) in acetic acid (50 mL) by reducing with sodium cyanoborohydride (1.3 g) in a manner similar to Example 88. The product (0.46 g) was purified by silica gel chromatography (5% methanol in EtOAc).

Example 89

20

8-Dimethylamino-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione hydrochloride



25

A solution of 2,5-di-(N²-dimethyl)amino-6-bromo-4-nitroindane (0.5 g, 1.7 mmol) and 20% Pd/C (0.1 g) in methanol (75 mL) was shaken on a Parr hydrogenation apparatus under a hydrogen atmosphere (52 psi) for 14.5 h. After removing the catalyst, the solvent was evaporated to give a white solid. The solid was dissolved in 2 N HCl (25 mL) and then treated with oxalic acid (0.5 g, 4 mmol) and heated at 100°C for 3

30

- 99 -


h. After cooling, the solution was basified, to pH 9 with 2 N NaOH. The resulting precipitate was collected by filtration and dried (0.4 g).

5

Example 90

8-Dimethylamino-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione

10



A mixture of 2,5-diamino-6-bromo-4-nitroindane (1 mmol) and paraformaldehyde (4 mmol) in acetic acid (100 mL) was treated portionwise with sodium cyanoborohydride (2 mmol), and then heated at 50°C for 4 h. The dimethylamino derivative was purified by column chromatography and converted by sequential hydrogenation and condensation with oxalic acid to the quinoxalinedione. Nitration at C-6 was accomplished with fuming nitric acid in trifluoroacetic acid at 0°C and then warming to room temperature. After stripping the solvent, the product was triturated with acetone, collected by filtration and washed with diethyl ether and dried in vacuo. Calc'd for $C_{17}H_{14}N_4O_4 \cdot HCl$: C, 47.79; H, 4.63; N, 17.15; found: C, 43.09; H, 4.08; N, 19.03. Ms M+1 (291).

30

Example 91

8-Diethylamino-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione hydrochloride

35

The title product was prepared in an identical manner to Example 89 using 2,5-di-(N²-diethyl)amino-6-bromo-4-nitroindane (0.46 g, 1.4 mmol) as a starting material.

- 100 -

Example 92

2-Ethylcarbamoylindane



A mixture of 2-aminoindane hydrochloride (25 g, 0.15 mol) and ethyl chloroformate (21.7 g, 0.2 mol) in saturated sodium bicarbonate (200 mL) and ether (200 mL) was stirred at room temperature for 1 h. The ether layer was separated and the aqueous layer back extracted. The combined ether extracts were washed with saturated NaCl, dried over magnesium sulfate, filtered and evaporated to give a white solid (27 g).

15

Example 93

2-(N-Methyl)acetamidoindane


20




A solution of 2-ethylcarbamoylindane (20.5 g, 0.1 mol) in diethyl ether (250 mL) was cooled in an ice bath and then treated portionwise with lithium aluminum hydride (7.6 g, 0.2 mol). The reaction mixture was stirred over 16 h while warming to room temperature. The reduction was incomplete, and was then heated at reflux for an additional 24 h. The mixture was cooled to 0°C and treated cautiously with water (8 mL), followed by 12.5% NaOH (6.5 mL) and water (16 mL). The white precipitate was removed by filtration and washed with ether and THF. The filtrate was treated with acetic anhydride (13 g) and the filtrate evaporated to a syrup. Silica gel chromatography (1:1 hexane:EtOAc) gave the product (12.2 g).

Example 94

2- (N-Methyl)acetamido-5-nitroindane



A mixture of 2- (N-methyl)acetamidoindane (16 g, 85 mmol) in trifluoroacetic acid (200 mL) was cooled in an ice bath and then treated with fuming nitric acid (20 mL). After stirring for 1 h in the ice bath, the reaction was stirred for an additional 3 h at room temperature. The solvent was removed and the residue taken up in methylene chloride:water. The aqueous phase was extracted with methylene chloride and the combined organic layers were dried over magnesium sulfate, filtered and evaporated to give the product (20 g).

Example 95N²-Methyl-2,5-diacetamidoindane

25 A mixture of 2- (N-methyl)acetamido-5-nitroindane (20.1 g, 85 mmol) and Raney nickel (10 g) in THF (100 mL) and methanol (100 mL) was stirred under a hydrogen atmosphere (balloon) at room temperature for 18 h. The catalyst was removed, and the filtrate was treated with acetic anhydride (18 g), and then evaporated to a brown syrup (22 g).

30

Example 96N²-Methyl-2,6-diacetamido-5-bromoindane

5 A mixture of N²-methyl-2,5-diacetamidoindane (22 g, 85 mmol) and sodium acetate trihydrate (23 g, 0.17 mmol) in acetic acid (250 mL) was treated dropwise with
10 bromine (17 g, 0.11 mol) and stirred for 18 h at room temperature. The mixture was treated with sodium bisulfite and evaporated. The residue was taken up in methylene chloride:water, and the aqueous layer was
15 back extracted with additional methylene chloride. The combined organic layers were dried over sodium sulfate, filtered and evaporated to give a syrup (26 g). The syrup was triturated with diethyl ether to give the product as a white solid.

Example 97N²-Methyl-2,5-diacetamido-6-bromo-4-nitroindane


25 A mixture of N²-methyl-2,6-diacetamido-5-bromoindane (21.8 g, 67 mmol) in trifluoroacetic acid (200 mL) was cooled in an ice bath and treated with fuming nitric acid (25 mL). After stirring for 1 h, the mixture was
30 warmed to room temperature and stirred for an additional 3 h. The solvent was removed and the residue taken up in water. The mixture was triturated in diethyl ether to give a precipitate, which was

collected by filtration and dried to give the product (22.5 g).

Example 98

5

N²-Methyl-2,5-diamino-6-bromo-4-nitroindane hydrochloride



10


A mixture of N²-methyl-2,5-diacetamido-6-bromo-4-nitroindane (22.5 g, 61 mmol) in 3 N HCl (150 mL) and acetic acid (50 mL) was heated at 100°C for 24 h. The mixture was cooled in an ice bath and the orange precipitate was collected by filtration and dried (17.2 g).

15

Example 99

20

8-Methylamino-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione



25

A mixture of N²-methyl-2,5-diamino-6-bromo-4-nitroindane hydrochloride (17.2 g, 53 mmol) and 20% Pd/C (1 g) in methanol (250 mL) was shaken on a Parr hydrogenation apparatus under a hydrogen atmosphere (50 psi) for 6 h. After removing the catalyst, the filtrate was evaporated and the residue taken up in 2N HCl (250 mL) and oxalic acid (13 g, 0.1 mmol) and heated at 100°C for 5 h. The mixture was cooled in a refrigerator

30

- 104 -


overnight, and the resulting precipitate was collected by filtration and washed with water and then diethyl ether and dried to give the product (9.9g).

5

Example 100

8-Methylamino-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione methane sulfonate salt

10



A mixture of 8-methylamino-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione (9.9 g) in trifluoroacetic acid was cooled in an ice bath and treated with fuming nitric acid (10 mL). After stirring at 0°C for 1 h, the reaction mixture was warmed to room temperature for an additional 1 h. The solvent was removed, and the residue taken up in acetone:water. The yellow solid was collected by filtration, washed with water and dried (9.8 g). One gram was taken up in water and NaOH (1 eq) and stirred at room temperature for 3 h. The solid was collected by filtration and dried. The solid was suspended in methanol and treated with methanesulfonic acid and stirred for 15 min. A tan solid was collected by filtration and dried.

8-Methylamino-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione. A mixture of 8-dimethylamino-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione hydrochloride (0.39 g) in trifluoroacetic acid was cooled in an ice bath and then treated with fuming nitric acid (1 mL). After

- 105 -

stirring for 1 h, the mixture was warmed to room temperature for 3 h. The solvent was evaporated and the residue suspended in methanol. The resulting yellow solid was collected by filtration and dried.

- 5 Calc'd for $C_{12}H_{12}N_4O_4 \cdot CH_3O_3S$: C, 41.93; H, 4.33; N, 15.05; found: C, 39.56; H, 4.11; N, 13.61. Ms M+1 (277).

Example 101

- 10 2-[Bis-(2-hydroxy-ethyl)-amino]-N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide

- A mixture of 2-[bis-(2-hydroxy)ethylglycine (1.6 g, 10 mmol), t-butyldimethylsilylchloride (6 g, 40 mmol) and imidazole (3.4 g, 50 mmol) in DMF (30 mL) was stirred at room temperature for 18 h. The solvent was evaporated and the residue dissolved in 1 N HCl:methylene chloride. The layers were separated, and the aqueous layer back extracted. The combined organic layers were dried over sodium sulfate, filtered and evaporated to give a solid. The resulting silylated derivative was suspended in hexane, collected by filtration and dried.

- 25 A solution of bis(2-t-butyldimethylsilyloxy)ethylglycine (0.38 g, 1 mmol) and carbonyldiimidazole (0.16 g, 1 mmol) in THF (5 mL) was heated at reflux for 30 min. The mixture was then added to a solution of 8-amino-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione hydrochloride (0.15 g, 0.5 mmol) in DMF (20 mL) and stirred at 70°C. After 30 min, triethylamine (0.2 g, 4 eq) was added and the reaction mixture was heated for 18 h. Another equivalent of the adduct of the starting glycine adduct and CDI was prepared as before, and added to the mixture. After heating at 70°C for 18 h longer, the mixture was filtered and the filtrate was evaporated. The residue was warmed in ethanol to give a solid,


- 106 -

which was collected and dried. Calc'd for $C_{17}H_{21}N_3O_7 \cdot HCl \cdot 0.4 \cdot H_2O$: C, 45.08; H, 5.12; N, 15.47; found: C, 44.71; H, 4.70; N, 15.65.

5

Example 102

5-Amino-6-bromo-4-nitro-2-pyrrolidin-1-yl-indane



10

A mixture of 2,5-diamino-6-bromo-4-nitroindane hydrochloride (1.5 g, 5 mmol), 1,4-dibromobutane (1.3 g, 6 mmol) and diisopropylethylamine (1.9 g, 15 mmol) in DMF (100 mL) was heated at 70°C for 24 h. The solvent was evaporated and the residue taken up in methylene chloride:water. The aqueous layer was basified with 1 N NaOH and back extracted. The combined organic layers were dried over sodium sulfate, filtered and evaporated to give a dark syrup. The purified product (0.5 g) was obtained from column chromatography (4:1 EtOAc:methanol).


15

20

Example 103

25

8-Pyrrolidin-1-yl-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione



30

The title product was obtained by an analogous procedure to that described in Example 100 employing

- 107 -

hydrogenation followed by condensation with oxalic acid to give the product (0.15 g).

Example 104

5

6-Nitro-8-pyrrolidin-1-yl-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione

10 Nitration of 8-pyrrolidin-1-yl-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione as described before with fuming nitric acid in trifluoroacetic acid gave the product as a yellow solid (0.1 g). $C_{13}H_{16}N_4O_4 \cdot HCl$. MS M+1 (317); HPLC 99%.

15

Example 105

20 N-Methyl-N-(6-nitro-2,3,-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-yl)-acetamide

A mixture of 8-methylamino-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione (0.5 g, 1.6 mmol), acetic anhydride (0.5 g, 5 mmol), triethylamine (0.5 g, 5 mmol) in water (5 mL) was stirred for 15 min at room temperature. The solid was collected by filtration, washed with 2 N HCl, then washed with diethyl ether and dried (0.5 g). Calc'd for $C_{14}H_{14}N_4O_3 \cdot C$, 52.83; H, 4.43; N, 17.60; found: C, 48.62; H, 4.26; N, 15.50.

30

Example 106

5-Amino-6-bromo-4-nitro-2-piperidin-1-yl-indane

35

A mixture of 2,5-diamino-6-bromo-4-nitroindane hydrochloride (1.5 g, 5 mmol), 1,5-dibromopentane (1.4 g, 6 mmol) and diisopropylethylamine (1.3 g, 10 mmol) in DMF (100 mL) was heated at 50°C for 24 h. The solvent was evaporated and the residue taken up in methylene chloride:water. The aqueous layer was

40

- 108 -

basified with 2 N NaOH and back extracted. The combined organic layers were dried over sodium sulfate, filtered and evaporated to give a syrup. The purified product (0.8 g) was obtained from column chromatography (4:1 EtOAc:methanol).

Example 107

8-Piperidin-1-yl-4,7,8,9-tetrahydro-
1H-cyclopenta[f]quinoxaline-2,3-dione
hydrochloride

The product (0.35 g) was obtained using a standard hydrogenation of 5-amino-6-bromo-4-nitro-2-piperidin-1-yl-indane (0.8 g, 2.6 mmol) over 20% Pd/C and condensation with oxalic acid.

Example 108

6-Nitro-8-piperidin-1-yl-4,7,8,9-tetrahydro-
1H-cyclopenta[f]quinoxaline-2,3-dione

A yellow solid was obtained by an identical procedure as used for example 61. Calc'd for $C_{16}H_{18}N_4O_4 \cdot HCl$: C, 52.39; H, 5.22; N, 15.27; found: C, 39.21; H, 4.60; N, 15.65 Ms M+1 (331).

Example 109

8-Diethylamino-6-nitro-4,7,8,9-tetrahydro-
1H-cyclopenta[f]quinoxaline-2,3-dione


Prepared by an identical procedure as used for example 61. Calc'd for $C_{18}H_{22}N_4O_4 \cdot HCl$: C, 42.39; H, 4.56; N, 16.29; found: C, 50.78; H, 5.40; N, 15.79. MS M+1 (319).

Other variations and modifications of this invention will be obvious to those skilled in the art.

- 109 -

WHAT IS CLAIMED IS:

1. A compound represented by the formula:



or a pharmaceutically acceptable salt thereof,
wherein

Z is an alicyclic fused ring having 5 to 7 carbon atoms;

R¹ is hydrogen, an alkyl or an arylalkyl;

X and Y are independently hydrogen, halogen, nitro, cyano, COOH, CONR²R³, SONR²R³ wherein R² and R³ are independently hydrogen, alkyl having 1 to 6 carbon atoms, cycloalkyl or aralkyl; and

A is O, NR⁴, CH₂NR⁴, CN, tetrazole or CO wherein R⁴ is hydrogen, alkyl, alkanol, alkyl amine or aralkyl, wherein

(i) when A is O, NR⁴ or CH₂NR⁴ then B is hydrogen, alkyl, alkenyl, alkynyl, aryl, aralkyl, alkanol, alkoxy, alkyl amine, heterocyclic, alkylheterocyclic, alkylcarbonyl, cycloalkylcarbonyl, arylcarbonyl, aralkylcarbonyl, heterocyclic-carbonyl, alkylheterocyclic-carbonyl, any of which may be unsubstituted or substituted by one or more hydroxy, thiol, amine, alkyl or butoxycarbonyl groups, CONR⁵R⁶ wherein R⁵ is hydrogen, alkyl having 1 to 6 carbon atoms, aralkyl or NR⁵ is a cyclic amine and R⁶ is alkyl, aryl, aralkyl or not present when NR⁵ is the cyclic amine, or when A is NR⁴ or CH₂NR⁴ then B is a naturally occurring α -amino acid moiety joined by an amide bond or B joins with R⁴ to form a four to seven membered heterocyclic ring, provided that when Z is a fused cyclohexyl ring and R⁴ is hydrogen then B is not hydrogen;

(ii) when A is CN then B is not present and Z is not a fused cyclohexyl ring;

(iii) when A is tetrazole then B is hydrogen or alkyl having 1 to 6 carbon atoms; and


(iv) when A is CO then B is hydroxy, alkoxy, aralkoxy, alkyl having 1 to 6 carbon atoms, aralkyl, NR^7R^8 wherein R^7 is hydrogen, alkyl having 1 to 6 carbon atoms, aralkyl or NR^7 is a cyclic amine and R^8 is alkyl, aryl, aralkyl or not present when NR^7 is the cyclic amine.

2. A compound according to claim 1, wherein Z is a cyclopentyl fused ring or a cyclohexyl fused ring, R^1 is hydrogen, X and Y are independently hydrogen, nitro or halogen, A is O, NR^4 , CH_2NR^4 , CN, tetrazole or CO, and R^4 is hydrogen, methyl, ethyl or joined with B to form a heterocyclic ring.

3. A compound according to claim 2, wherein B is hydrogen, alkyl, aryl, aralkyl, heterocyclic, alkylheterocyclic, alkylcarbonyl, cycloalkylcarbonyl, arylcarbonyl, aralkylcarbonyl, heterocyclic-carbonyl, alkylheterocyclic-carbonyl, or alkoxy.


4. A compound according to claim 3, wherein A is O, NR^4 or CO and X is nitro.

5. A compound according to claim 4, wherein Z is a cyclohexyl fused ring, A is O, NR^4 or CH_2NR^4 and B is



- 111 -

6. A compound according to claim 4, wherein Z is a cyclopentyl fused ring, A is NR^4 or CH_2NR^4 , and B is methyl, ethyl, acetyl,



7. A compound according to claim 2, wherein the heterocyclic ring is a pyrrolidine ring or a piperidine ring.

- 112 -

8. A compound selected from the group consisting of:

9-benzyloxy-6-bromo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

6-bromo-9-hydroxy-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

piperidine-4-carboxylic acid (6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-octahydro-benzo[f]quinoxalin-9-yl) amide;

N-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-octahydro-benzo[f]quinoxalin-9-yl)-acetamide;

9-benzylamino-6-bromo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

N-(5-bromo-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide;

N-(5-bromo-6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide;

N-(2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide;

N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide;

8-amino-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione hydrochloride;

4-(2,3,-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-ylcarbamoyl)-piperidine-1-carboxylic acid tert-butyl ester;

- 113 -

4-(6-nitro-2,3,-dioxo-2,3,4,7,8,9-hexahydro-1H
cyclopenta[f]quinoxalin-8-yl-carbamoyl)-piperidine
hydrochloride;

N-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-
octahydrobenzo[f]quinoxalin-9-yl)-benzamide;

N-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-
octahydrobenzo[f]quinoxalin-9-yl) cyclohexylamide;

8-amino-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]-
quinoxaline-2,3-dione hydrochloride;

N-(6-nitro-2,3-dioxo-4,7,8,9-tetrahydro-1H
cyclopenta[f]quinoxalin-8-yl)-benzamide;

4-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-
octahydrobenzo[f]quinoxalin-9-ylcarbamoyl)-piperidine
hydrochloride;

2-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-
octahydrobenzo[f]quinoxalin-9-ylcarbamoyl) ethyl-(4-
hydroxy) benzene;

N-phenyl-N'-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10 -
octahydrobenzo[f]quinoxalin-9-yl)-urea;

N-methyl-N-(6-nitro-2,3-dioxo-1,2,3,4,7,8,9,10-
octahydro-benzo[f]quinoxalin-9-yl)-benzamide;

thiophene-2-carboxylic acid (6-nitro-2,3-dioxo-
2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-
yl)-amide;

furan-2-carboxylic acid (6-nitro-2,3-dioxo-2,3,4,7,8,9-
hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-amide;

thiophene-3-carboxylic acid (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)amide;

N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-nicotinamide;

pyridine-2-carboxylic acid (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-amide;

2,3,4,5-tetrahydroxy-tetrahydro-furan-2-carboxylic acid (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-amide;

pyrrolidine-2-carboxylic acid (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-amide;

benzo[b]thiophene-2-carboxylic acid (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-amide;

2-(1H-indol-3-yl)-N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide;

thiophene-2-carboxylic acid methyl-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-amide;

N-methyl-N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-benzamide;

6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-carboxylic acid methyl-phenyl-amide;

- 115 -

6-nitro-2,3,-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-carboxylic acid;

6-nitro-2,3,-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-carboxylic acid phenylamide;

9-methylamino-6-nitro-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

6-nitro-9-pyrrolidin-1-yl-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

8-dimethylamino-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

8-methylamino-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione methane sulfonate salt;

2-[bis-(2-hydroxy-ethyl)-amino]-N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide;

6-nitro-8-pyrrolidin-1-yl-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

N-methyl-N-(6-nitro-2,3,-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxaline-8-yl)-acetamide;

6-nitro-8-piperidin-1-yl-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

8-diethylamino-6-nitro-4,7,8,9,-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

6-nitro-8-(thiophen-2-ylmethoxy)-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

6-nitro-9-(thiophen-2-ylmethoxy)-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

6-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yloxymethyl)-1,2,3,4-tetrahydro-isoquinoline-3-carboxylic acid;

6-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yloxymethyl)-1,2,3,4-tetrahydro-isoquinoline-3-carboxylic acid;

6-nitro-8-[2-(1H-pyrrol-2-yl)-ethoxy]-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

6-nitro-9-[2-(1H-pyrrol-2-yl)-ethoxy]-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

6-nitro-8-[[1H-tetrazol-5-ylmethyl)-amino]-methyl]-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

6-nitro-9-[[1H-tetrazol-5-ylmethyl)-amino]-methyl]-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

6-nitro-8-[[1H-tetrazol-5-ylmethyl)-amino]-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

6-nitro-8-[[1H-tetrazol-5-ylmethyl)-amino]-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

8-[[methyl-(1H-tetrazol-5-ylmethyl)-amino]-methyl]-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

- 117 -

9-({methyl-(1H-tetrazol-5-ylmethyl)-amino}-methyl)-6-nitro-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

6-nitro-8-[2-(1H-tetrazol-5-yl)-ethoxy]-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

6-nitro-9-[2-(1H-tetrazol-5-yl)-ethoxy]-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

6-nitro-8-[2-(1H-tetrazol-5-yl)-ethylamino]-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

6-nitro-9-[2-(1H-tetrazol-5-yl)-ethylamino]-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

6-nitro-8-(1H-tetrazol-5-ylmethoxy)-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

6-nitro-9-(1H-tetrazol-5-ylmethoxy)-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-ylmethyl)-2-(1H-tetrazol-5-yl)-acetamide;

N-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-ylmethyl)-2-(1H-tetrazol-5-yl)-acetamide;

2-(3-hydroxy-5-methyl-isoxazol-4-yl)-N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide;

- 118 -

2-(3-hydroxy-5-methyl-isoxazol-4-yl)-N-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl)-acetamide;

2-(3-hydroxy-5-methyl-isoxazol-4-yl)-N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-ylmethyl)-acetamide;

2-(3-hydroxy-5-methyl-isoxazol-4-yl)-N-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-ylmethyl)-acetamide;

8-(3-hydroxy-5-methyl-isoxazol-4-ylmethoxy)-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

9-(3-hydroxy-5-methyl-isoxazol-4-ylmethoxy)-6-nitro-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-2-(1H-tetrazol-5-yl)-acetamide;

N-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl)-2-(1H-tetrazol-5-yl)-acetamide;

3-[(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-ylmethyl)-carbamoyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid;

3-[(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-ylmethyl)-carbamoyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid;

- 119 -

8-[2-(3-hydroxy-5-methyl-isoxazol-4-yl)-ethoxy]-6-nitro-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

9-[2-(3-hydroxy-5-methyl-isoxazol-4-yl)-ethoxy]-6-nitro-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

2-(2-hydroxy-phenyl)-N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-acetamide;

2-(2-hydroxy-phenyl)-N-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl)-acetamide;

2-hydroxy-N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-ylmethyl)-benzamide;

2-hydroxy-N-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-ylmethyl)-benzamide;

6-nitro-8-(2-pyridin-2-yl-ethoxy)-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

6-nitro-9-(2-pyridin-2-yl-ethoxy)-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-2-[2-(1H-tetrazol-5-yl)-phenyl]-acetamide;

N-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl)-2-[2-(1H-tetrazol-5-yl)-phenyl]-acetamide;

- 120 -

1-methyl-2,4-dioxo-1,2,3,4-tetrahydro-pyrimidine-5-carboxylic acid (6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-ylmethyl)-amide;

1-methyl-2,4-dioxo-1,2,3,4-tetrahydro-pyrimidine-5-carboxylic acid (6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-ylmethyl)-amide;

6-nitro-8-(5-oxo-4,5-dihydro-[1,3,4]oxadiazol-2-ylmethoxy)-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

6-nitro-9-(5-oxo-4,5-dihydro-[1,3,4]oxadiazol-2-ylmethoxy)-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

6-nitro-8-(pyridin-2-ylamino)-4,7,8,9-tetrahydro-1H-cyclopenta[f]quinoxaline-2,3-dione;

6-nitro-9-(pyridin-2-ylamino)-1,4,7,8,9,10-hexahydro-benzo[f]quinoxaline-2,3-dione;

4-{{{6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-ylmethyl)-amino}-methyl}-benzoic acid;

4-{{{6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-ylmethyl)-amino}-methyl}-benzoic acid;

2-hydroxy-benzoic acid 6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl ester;

2-hydroxy-benzoic acid 6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl ester;

- 121 -

N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-phthalamic acid;

N-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl)-phthalamic acid;

2-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yloxy)-benzoic acid;

2-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl oxy)-benzoic acid;

(3-hydroxy-5-methyl-isoxazol-4-yl)-acetic acid 6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl ester;

(3-hydroxy-5-methyl-isoxazol-4-yl)-acetic acid 6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl ester;

2-hydroxy-N-(6-nitro-2,3-dioxo-2,3,4,7,8,9-hexahydro-1H-cyclopenta[f]quinoxalin-8-yl)-benzamide;

2-hydroxy-N-(6-nitro-2,3-dioxo-1,4,7,8,9,10-hexahydro-benzo[f]quinoxalin-9-yl)-benzamide; and

a pharmaceutically acceptable salt thereof.

9. A pharmaceutical composition comprising a pharmaceutically acceptable carrier or diluent and a therapeutic amount of at least one compound of claim 1.

10. A method of therapeutic treatment comprising administering to a human at least one compound of claim 1 in unit dosage form.

- 122 -

11. A method for treating anxiety which comprises administering at least one compound of claim 1 in unit dosage form.


12. A method for treating cerebral hypoxic/ischemia which comprises administering at least one compound of claim 1 in unit dosage form.

13. A method for treating Parkinsonism which comprises administering at least one compound of claim 1 in unit dosage form.


14. A method for treating a human with an anticonvulsant comprising administering to a human at least one compound of claim 1 in unit dosage form.

15. A method for preparing the compound of formula (I) comprising the steps of:

(a) reacting a compound of formula




wherein Z is an alicyclic fused ring having 5 to 7 carbon atoms and A is OH, N₃, COOH, COO alkyl, COO-benzyl, a heterocyclic ring or a protected amine, with a halogenating reagent to produce a compound of formula




wherein X is a halogen,

- 123 -

(b) nitrating the compound formed in step (a) to produce a compound of formula




(c) sequentially hydrolyzing an acetamide of the compound of step (b) and then catalytically hydrogenating the hydrolyzed compound to produce a compound of formula




wherein X' is a halogen or hydrogen,


(d) condensation of the compound of step (c) with oxalic acid or an oxalic acid ester to produce a compound of formula



(e) optionally hydrogenating the compound of step (d) when X' is halogen to produce a compound of formula




(f) optionally reacting the compound of steps (d) or (e) with an electrophilic substituent to produce a compound of formula



wherein X and Y are independently hydrogen, halogen, nitro, cyano, COOH, CONR²R³, SONR²R³, wherein R² and R³ are independently hydrogen, alkyl, cycloalkyl or aralkyl; and

(g) optionally reacting the compound of steps (d), (e) or (f) when A' is amine or OH with a reagent selected from the group consisting of electrophilic reagents and hydride reagents to produce a compound of formula



wherein A and B are as defined for formula (I) in claim 1.

16. A method according to claim 15, wherein the halogenating agent in step (a) is selected from the group consisting of bromine, N-bromosuccinimide, chlorine, and iodochloride.

17. A method according to claim 15, wherein the compound formed in step (a) is nitrated in step (b) with fuming nitric acid or potassium nitrate in an acid

- 125 -


selected from the group consisting of trifluoroacetic acid, acetic acid or concentrated sulfuric acid.

18. A method according to claim 15, wherein the hydrogenation step in steps (c) or (d) comprises hydrogenation with a nickel or palladium catalyst in a hydrogen atmosphere.


19. A method according to claim 15, wherein the electrophilic reagents of step (g) are selected from the group consisting of alkyl halides, acylhalide, aldehydes, isocyanates and mixtures thereof.

20. A method for preparing substituted cyclopentyl fused ring quinoxalinediones comprising the steps of:


(a) acetylating and nitrating a compound of the formula



to form a compound of the formula




(b) hydrogenation of a nitro group, acetylating a resulting aniline and halogenating the compound formed in step (a) to form a compound of the formula




wherein X is a halogen,

(c) nitrating the compound of step (b) to form a compound of the formula


- 126 -



(d) hydrolysis of the compound of step (c) to form a compound of the formula




(e) hydrogenating the compound of step (d) to form a diamine compound of the formula



or


reacting the compound of step (d) with XCCOOR^{16} ,

wherein R^{16} is hydrogen, alkyl or aralkyl, to form an amide compound of the formula




(f) optionally hydrogenating the diamino compound of step (d) to form a compound of the formula

- 127 -




(g) condensation of the diamino compound of step (d) or the compound of step (f) with oxalic acid or an oxalic acid ester or hydrogenation of the amide compound of step (e) to form a compound of the formula




wherein X' is a halogen or hydrogen, or

(h) optionally reacting the compound of step (g) with an electrophilic substituent to produce a compound of formula



wherein X and Y are the same as defined for formula (I) in claim 1, and

(i) optionally hydrolyzing compound of steps (g) or (i) with a reagent selected from the group consisting of electrophilic reagents and hydride reagents to produce a compound of the formula



wherein R is hydrogen, alkyl, alkenyl, alkynyl, aryl, aralkyl, alkylheterocyclic, alkylcarbonyl, aralkylcarbonyl, alkylheterocycliccarbonyl, CONR⁵R⁶ wherein R⁵ is hydrogen, alkyl, aralkyl or NR⁵ is a cyclic amine and R⁶ is alkyl, aryl, aralkyl or not present, or a common amino acid moiety joined by an amide bond.

21. A method according to claim 20, wherein in step (a) the step of acetylating is performed with acetyl chloride or acetic anhydride.

22. A method according to claim 20, wherein in steps (b), (e) or (f) the step of hydrogenating is performed with a nickel or palladium catalyst in a hydrogen atmosphere.

23. A method according to claim 20, wherein in step (b) the step of halogenating is performed with a halogenating agent selected from the group consisting of chlorine, bromine or iodine monochloride.

24. A method according to claim 20, wherein in steps (a) or (c) the step of nitrating is performed with fuming nitric acid or potassium nitrate in an acid selected from the group consisting of acetic acid, trifluoroacetic acid or concentrated sulfuric acid.


25. A method according to claim 20, wherein the electrophilic reagents of step (i) are selected from

- 129 -


the group consisting of alkyl halides, acylhalide, aldehydes, isocyanates and mixtures thereof.

26. A method for preparing the compound of formula (I) comprising the steps of:


(a) reacting a compound of formula




wherein Z is an alicyclic fused ring having 5 to 7 carbon atoms and A is OH, N₃, COOH, COO-alkyl, COO-benzyl, a heterocyclic ring or a protected amine, with a brominating agent to produce a compound of formula



(b) nitrating the compound formed in step (a) to produce a compound of formula




(c) catalytically hydrogenating the compound of step (b) to produce a compound of formula




- 130 -


(d) condensation of the compound of step (c) with oxalic acid or an oxalic acid ester to produce a compound of formula



(e) optionally hydrogenating the compound of step (d) to produce a compound of formula




(f) optionally reacting the compound of steps (d) or (e) with an electrophilic substituent to produce a compound of formula



wherein X and Y are independently hydrogen, halogen, nitro, cyano, COOH, CONR²R³, SONR²R³, wherein R² and R³ are independently hydrogen, alkyl, cycloalkyl or aralkyl; and

(g) optionally reacting the compound of steps (d), (e) or (f) when A' is NR⁴H or OH with a reagent selected from the group consisting of electrophilic reagents and hydride reagents to produce a compound of formula


- 131 -




wherein A, B and R⁴ are as defined for formula (I) in claim 1.

27. A method for preparing substituted cyclopentyl fused ring quinoxalinediones comprising the steps of:


(a) acetylating and nitrating a compound of the formula




to form a compound of the formula



(b) reducing, acetylating and brominating the compound formed in step (a) to form a compound of the formula




(c) nitrating the compound of step (b) to form a compound of the formula




- 132 -


(d) hydrogenating the compound of step (c) to form a compound of the formula




(e) optionally hydrogenating the compound of step (d) to form a compound of the formula



(f) condensation of the compound of steps (d) or (e) with oxalic acid or an oxalic acid ester to form a compound of the formula




(g) optionally reacting the compound of step (f) with an electrophilic substituent to produce a compound of formula



wherein X and Y are the same as defined for formula (I) in claim 1, and



- 133 -

(h) optionally reacting the compound of steps (f) or (g) with a reagent selected from the group consisting of electrophilic reagents and hydride reagents to produce a compound of the formula




wherein R is hydrogen, alkyl, alkenyl, alkynyl, aryl, aralkyl, alkylheterocyclic, alkylcarbonyl, aralkylcarbonyl, alkylheterocycliccarbonyl, CONR^5R^6 wherein R^5 is hydrogen, alkyl, aralkyl or NR^5 is a cyclic amine and R^6 is alkyl, aryl, aralkyl or not present, or a common amino acid moiety joined by an amide bond.

28. A compound selected from the group consisting of the compounds by the formulae II, III, IV and V:



wherein R^9 is N , or $\text{R}^{10}\text{HN}-$ wherein R^{10} is hydrogen alkyl, alkenyl, alkynyl, aryl, aralkyl, alkylheterocyclic, alkylcarbonyl, aralkylcarbonyl, alkylheterocycliccarbonyl, CONR^5R^6 wherein R^5 is hydrogen, alkyl, aralkyl or NR^5 is a cyclic amine and R^6 is alkyl, aralkyl, or a common amino acid moiety joined by an amide bond,

- 134 -




(IV)

wherein A' is OH, N₃ or R¹⁰NH— and R¹⁰ is as previously described, Z is an alicyclic fused ring having 5 to 7 carbon atoms, R¹¹ and R¹² are independently NHAc,

NHCCOOR^{17} , NH_2 , NHCOCOOR^{15} wherein R¹⁵ is hydrogen or

alkyl, or NO₂, and R¹³ and R¹⁴ are independently hydrogen or Br provided that at least one of R¹³ and R¹⁴ is Br, R¹⁷ is hydrogen alkyl or aryl, and



(V)

wherein A' and Z are as previously described provided that when Z is a fused cyclohexyl ring R¹⁰ is not hydrogen, and wherein R¹, X and Y are as described for formula (I) in claim 1.

29. A compound according to claim 28, wherein Z is a fused cyclohexyl ring, R¹¹ is NHAc, NHCOCOOR^{15} or NH_2 , R¹² is NO₂, or NH_2 , R¹³ is hydrogen and R¹⁴ is Br.

30. A compound according to claim 28, wherein Z is a fused cyclopentyl ring, R¹¹ is NO₂, or NH_2 , R¹² is NHAc, NHCOCOOR^{15} or NH_2 , R¹³ is Br and R¹⁴ is hydrogen.

- 135 -

31. A compound selected from the group consisting of:

acetic acid 8-acetylamino-5-bromo-7-nitro-1,2,3,4-tetrahydronaphthalen-2-yl ester;

8-amino-5-bromo-7-nitro-1,2,3,4-tetrahydronaphthalen-2-ol;

7,8-diamino-5-bromo-1,2,3,4-tetrahydronaphthalen-2-ol;

methanesulfonic acid 8-acetyl-1,2,3,4-tetrahydronaphthalen-2-yl ester;

N-(7-azido-5,6,7,8-tetrahydronaphthalene-1-yl)-acetamide;

N-(7-azido-4-bromo-5,6,7,8-tetrahydronaphthalen-1-yl)-acetamide;

N-(7-azido-4-bromo-2-nitro-5,6,7,8-tetrahydronaphthalen-1-yl) acetamide;

7-azido-4-bromo-2-5,6,7,8-tetrahydro-naphthalen-1-ylamine;

N-7-benzyl-4-bromo-2-nitro-5,6,7,8-tetrahydronaphthalene-1,7-diamine;

N-(5-acetylamino-6-bromo-indan-2-yl)-acetamide;

N-(5-acetylamino-6-bromo-4-nitro-indan-2-yl)-acetamide;

N-(5-amino-6-bromo-4-nitro-indan-2-yl)-acetamide;

- 136 -

6-bromo-4-nitro-indan-2,5-diamine monohydrochloride;
N-(4,5-diamino-6-bromo-indan-2-yl)-acetamide; and
a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

Int: Int. Application No

PCT/US 95/14571

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D241/38 C07D403/04 C07D401/12 C07C233/54 C07C247/14
C07C233/43 C07C233/56 A61K31/495

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| X | EP,A,0 283 959 (A/S FERROSAN) 28 September 1988 cited in the application see the whole document | 1,9 |
| P,X | US,A,5 389 687 (J.SCHAUS ET AL.) 14 February 1995 see column 1 - column 2; claims; table II | 28 |



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "A" document member of the same patent family

Date of the actual completion of the international search

19 March 1996

Date of mailing of the international search report

25. 03. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Francois, J

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 95/14571

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| EP-A-283959 | 28-09-88 | AU-B- 596737 | 10-05-90 |
| | | AU-B- 1326088 | 22-09-88 |
| | | CA-A- 1315789 | 06-04-93 |
| | | DE-A- 3867268 | 13-02-92 |
| | | FI-C- 92582 | 12-12-94 |
| | | FI-B- 92582 | 31-08-94 |
| | | IE-B- 60002 | 18-05-94 |
| | | NO-B- 174549 | 14-02-94 |
| | | US-A- 4889855 | 26-12-89 |
| | | US-A- 4912108 | 27-03-90 |
| | | ZA-A- 8801139 | 15-08-88 |
| ----- | | | |
| US-A-5389687 | 14-02-95 | CA-A- 1335106 | 04-04-95 |
| ----- | | | |

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.